

Deranged Levels of Renal Parameters after Using Heavy Metal Kushta of Copper in Diet of Albino Rats

HIRA KAREEM¹, NADIA NASEEM², A H NAGI³, ROMMAN UL HAQ⁴.

ABSTRACT

Aim: To evaluate the effects of copper preparations of Kushta on renal parameters. This study was conducted from 14th of May 2013 to 30th of November 2013 in Pathology Department of University of Health Sciences Lahore and it was an experimental study.

Method: Thirty albino rats were selected and divided into three groups. The control group A was fed on standard pellet rodent diet and tap water. The experimental group B was fed single dose of copper preparation of Kushta (0.15mg/kg of body weight) for 18 weeks while experimental group C was fed double dose of copper preparation of Kushta (0.30mg/kg of body weight) for 18 weeks. Renal parameters were calculated once at the start of experiment and then after every 6 weeks up to 18 weeks.

Results: Results showed that renal parameters in terms of serum creatinine level and proteinuria were deranged evident on biochemical analysis. When the Control group A on normal diet was compared with the experimental group B and C, statistically significant difference ($p < 0.05$) was noticed in terms of renal parameters.

Conclusion: It is therefore concluded that copper preparations of Kushta deranged renal parameters in terms of serum creatinine levels and proteinuria.

Keywords: Kushta, heavy metals, copper, renal toxicity.

INTRODUCTION

Copper is a trace element vital for all human beings¹. It is an essential requirement in the synthesis of chlorophyll and haemoglobin². Cu metal is found naturally in the Earth's crust. Its natural air and water discharges are therefore considerable³. Heavy metals are those metallic elements which are dangerous at higher concentration and have greater density than 5g/cm³. Heavy metals are toxic in nature since they tend to bioaccumulate in the body. Some adverse effects of heavy metal products can also result from metal poisoning but is dose dependant. Usually people inhale or ingest small amounts of metals which gradually interfere with body functions and pathogens which may cause eventually death⁴. In the Earth's crust, many metals are found as natural constituents but many human procedures have altered their geochemical cycles and biochemical balance. Some form of pharmacological activity for example results in the accretion of metals in certain parts of the plant in the form of secondary metabolites. Any contact beyond the upper tolerable range, with these heavy metals such as cadmium, copper, lead, nickel and zinc can cause lethal effects

¹Demonstrator Microbiology SIMS, Lahore.

²Prof. Pathology UHS, Lahore.

³Assistant Prof. Pathology UHS, Lahore.

⁴Consultant Radiologist, Govt. Hospital Shahdra Lahore

Correspondence to Dr. Hira Kareem Email:

romman01@yahoo.com

to the humans. Heavy metals interrupt the metabolic processes in two ways. Once they disturb the functions of essential organs such as the heart, brain, kidneys, bones, liver, etc., and secondly, they shift critical mineral nutrients from their position of origin, and thus impede their natural organic purpose. However, it is not possible to make the surroundings free from these metals. There are several ways for heavy metals to be introduced in the body for example food consumption, beverages, skin exposure, inhaled air, soils, manufacturing activity and the utilization of fuel from the remnants. A contact to heavy metals such as cadmium, copper, lead, nickel and zinc is reported to be dangerous for the human health.⁵ Ingredients of "Kushta" by quakes and so-called hakims for the treatment of chronic conditions can damage the kidneys and can cause renal failure. Heavy metals are used in the synthesis of Kushta by hakims. The metals used in the preparation of Kushta aren't useful for health or digestion. They accumulate within the kidneys, creating a continuous contamination that ends up in severe renal impairment. Copper exists in the blood in two forms: one is bound to ceruloplasmin (85-95%), and the second is the 'free' rest, loosely bound to albumin and small molecules. Copper is transported in the blood by a protein ceruloplasmin. After absorption, it is transported to the liver. After oral intake it accumulates within the liver initially when it reaches the lethal level, pathological reaction

starts. Copper is absorbed in the duodenum and upper parts of the small intestine. Then 95 percent Cu ions are bound to protein ceruloplasmin and rest is with albumin and amino acids. A small amount is excreted in the urine. The main part of the Cu from the body is excreted in bile. Traffic and a good use of copper in the body are dependent on the proper functioning of many organs such as liver, gall bladder and kidneys. If any of these organs are not functioning properly, the body cannot use and excrete Cu properly. Copper is a strong oxidant metal causing inflammation and free radical damage to tissues. To avoid these toxic effects, it should be linked to binding proteins, ceruloplasmin and metallothionein. With increase in retention of copper, it produces a toxic effect (similar to other heavy metals) on the body and contributes several chronic diseases.⁴ As the kidneys are highly sophisticated transport organs, they excrete metabolic waste. Such a function is optimised by a special circulatory structure in the kidney, which has a remarkable ability to adjust the haemodynamic inputs in order to maintain renal circulation⁶. The basic functional unit of kidney is the nephrons. About 1.2 million of nephrons are present in each kidney, urine is synthesized by them. Each nephron is composed of Bowman capsule that surrounds the glomerular capillary tuft, the proximal convoluted tubule, the loop of Henle and the distal convoluted tubule, which empties into the collecting ducts⁷. All heavy metals are deposited in kidney tissue. When liver is being saturated with copper and all binding sites are occupied then copper compounds will begin to accumulate in the kidney⁸.

Renal lesions after using copper preparations of Kushta can happen by three possibilities. It could be acting as haptens, could intervene in the body's defense mechanism and act as direct foreign toxic substances like other heavy metals.⁹ The toxicity of these preparations of heavy metals depend on various factors such as the toxicity of metals, the duration of the exhibition, the total absorbed dose, age of the exposed person and the route of exposure.¹⁰ Kidneys act as filtration unit of the body so when these copper products have been used by humans they are filtered and trapped in the renal tubules and glomeruli, causing damage to the kidneys. Sometimes, it could be irreversible damaged the kidney and can lead to death. Kidney damage in terms of glomerulonephritis would be occurred by the use of these toxic metallic Kushtas. Glomerulonephritis is usually associated with rapidly progressive glomerulonephritis syndrome, which can occur mostly in the form of inflammation of glomeruli, including post infectious glomerulonephritis, IgA, lupus, Vasculitis renal nephropathy, membrano-

proliferative glomerulonephritis, and anti-glomerular basement (GBM) antibodies which are mainly caused by the toxicity of heavy metals.

MATERIAL AND METHODS

It was an experimental intervention, randomized controlled study in adult rats. The study was conducted at University of Health Sciences, Lahore. The animals were kept in the animal house and experimental work was further processed in the pathology department of University of Health Sciences, Lahore. The total duration of this study was 18 weeks. Thirty male and female albino rats of Wister strain, 6-8 weeks of age; weighing 200-250 Gms were procured from University of Health Sciences Lahore. Albino rats were separated in different cages and maintained in the animal house of the University of Health Sciences Lahore under controlled environment (temperature 22-25 c, humidity 65%±5) and light and dark cycle of 12 hours each. Each rat was to receive the prescribed dose on alternate days. Cage cards were used to indicate the group of albino rats.

1) **Group A:** is the control group. It included 10 healthy rats receiving normal diet and tap water for 18 weeks

2) **Group B:** It included 10 rats; Cu preparations of Kushta 0.15 mg/kg body weight, mixed and homogenized with wheat flour, and dispensed as pellets, were given orally to the rats of this group on alternate days for 18 weeks.

3) **Group C:** It included 10 rats and Cu preparations of Kushta 0.30 mg/kg body weight, mixed and homogenized with wheat flour, and dispensed as pellets, was given orally to the rats of this group on alternate days for 18 weeks.

Experimental schedule: All rats were weighed before the commencement of the experiment. No sex differences were made and all rats were kept in separate cages. They received nutritionally standard diet and water. After acclimatization, rats were divided into three groups as mentioned above. Each rat was marked for its identification by giving a number on its back. Before the start of experiment urinary proteins and serum creatinine level were measured in all rats.

Copper preparations of kushta: Copper preparations of Kushta used in this study were analyzed chemically from Biochemistry Department of University of Health Sciences Lahore, for exact quantification of the heavy metal. According to the body weight (0.5mg/kg body weight) doses were calculated, mixed and homogenized with wheat flour and dispensed as pallets. This dose has been used in various studies^{11,12}. Oral LD₅₀ of rats for Copper

Kushta has been reported as 2mg/kg body weight.¹³ Doses have been translated from human dose to rat dose by normalization of the body surface area (Animal dose mg/kg=Human Dose mg/kg multiplied by Animal Km/Human Km); where Km is 5.9 for rat and 37 for human¹⁴. The doses were adjusted daily according to the weight of each rat.

Methods for determination of serum creatinine and proteinuria: Serum creatinine was measured by using commercially available kits prepared by Randox (Ref: CR510, LOT: 216982). Urinary proteins were determined by strip method¹⁵.

Collection of blood sample: Method for obtaining blood sample and 24 hour urinary proteins: Blood samples from each group were collected by cardiac puncture.¹⁶ Albino rat was taken out by holding its tail and was anaesthetized by placing it in a plastic container having a tight lid covering and a cotton ball soaked in ether alcohol until the rat was completely anaesthetized. The rat was when removed from the container and 1 ml of blood was drawn in 3 ml disposable syringe by cardiac puncture. Blood was allowed to stand for one hour before centrifuging it at a speed of 3000r/pm for 10 minutes. Samples were collected in metabolic cages. During this collection of 24 hour urine samples the animals were allowed only water to drink.

RESULTS

Renal functions tests were performed once before starting the experiment as well as every six week thereafter for 18 weeks.

Table 1: Descriptive statistics for Creatinine level in all groups at different time intervals

Groups	N	Mean	SE
6 weeks of experiment			
Group-A	10	0.16	0.01
Group-B	10	0.48	0.03
Group-C	10	0.40	0.05
12weeks of experiment			
Group-A	10	0.16	0.01
Group-B	10	0.58	0.02
Group-C	10	0.54	0.02
18 weeks of experiment			
Group-A	10	0.16	0.01
Group-B	10	0.73	0.03
Group-C	10	0.73	0.11

Serum creatinine level: Serum creatinine level was assessed in all 3 groups of animals at the start of experiment and then after every 6 weeks. During this follow up time period it was observed that creatinine level in Group-A animals was 0.16±0.01 when assessed after 6, 12 and 18 weeks. While in Group-B & C creatinine level was quite high as compared to

Group-A rats at follow up time intervals, i.e., at 6, 12 and 18 weeks respectively.

24 hour urinary proteins level:: 24 hour Urinary proteins level in Group-A rats was 0.00 during the course of follow up time period i.e., (6,12 and 18 weeks). But 24 hour urinary proteins level in Group-B & C rats show high level of urea protein level at 6, 12 and at 18 weeks (Table 2).

Table 2: Descriptive statistics for 24 hour urinary Proteins level in all groups at different time intervals

Groups	N	Mean	SE
6 weeks of experiment			
Group-A	10	0.00	0.00
Group-B	10	142.00	60.73
Group-C	10	139.00	61.41
12weeks of experiment			
Group-A	10	0.00	0.000
Group-B	10	213.00	63.00
Group-C	10	173.00	54.93
18 weeks of experiment			
Group-A	10	0.00	0.00
Group-B	10	340.00	65.31
Group-C	10	380.00	61.10

DISCUSSION

The present study demonstrated that copper in dose dependent manner is renal toxicant and it changes serum parameter levels related to renal function. Serum levels of creatinine increased as a result of copper exposure in comparison to the control group. These observed changes are in accordance with other authors' results^{13,19}.

The significant role of kidneys in both excretory and endocrine process cannot be denied. Kushta and metal containing products hinder the function of kidneys.

The scope of the present study is to highlight the toxic effects of copper preparations of Kushta on kidneys of albino rats. In the present study rats, were treated with various doses of copper preparations of Kushta orally and markedly raised serum creatinine level were observed during the study except in control group. Six rats from group C and 10 rats from group D showed up to 1.1mg/dl. Elevated creatinine level signifies impair kidney function or kidney disease and this relation has been very well established in various other experimental conditions^{19,20}. Serum levels of creatinine increased as a result of copper exposure in comparison to the control group. These changes are in accordance with other authors' results^{17,18}.

Similarly in the present study significant association has been found between various doses of copper preparation of Kushta and proteinuria. One

of the most toxic effects of copper on kidney is proteinuria.

After six weeks four rats showed proteinuria of 30mg/dl and six rats showed proteinuria of 100mg/dl of group B while in group C, one rat showed 30mg/dl proteinuria and four rats showed 100mg/dl proteinuria and five rats showed 500mg/dl proteinuria. After twelve weeks nine rats from group B and three rats from group C showed 100 mg/dl proteinuria and one from group B and seven from group C showed 500mg/dl proteinuria. At the end of experiment eight rats from group B and all rats from group C showed 500 mg/dl proteinuria. This showed the clear association between doses of metallic Kushta and proteinuria which led to the irreversible kidney damage. These findings are consistent with the other studies describing indication of kidney damage is proteinuria²¹. One of the most toxic effects of copper on kidney is proteinuria²⁰.

The present study demonstrated that copper in a dose dependent manner is renal toxicant and it changes serum parameter levels related to renal function.

CONCLUSION

In conclusion, present study clearly suggests that copper in dose-dependent manner by changing serum parameters related to renal function can induce renal toxicity. It is therefore concluded that copper preparations of Kushta deranged renal parameters in terms of serum creatinine levels and proteinuria.

REFERENCES

1. deMayo A. and Taylor M.C. Guidelines for Surface Water Quality. Inorganic Chemical Substances. Copper. Water Quality Branch, Inland Waters Directorate, Environment Canada, Ottawa. (1981); (1).
2. Birge W J and Black J A. Effects of Copper on Embryonic and Juvenile Stages of Aquatic Animals. (1997); Nriagu JO ed. Copper in Environment. Part 2; 373-399. John Wiley NY.
3. Barceloux D G. Copper. J Toxicol Clinical Toxicology. (1999); 37(2):217-230.
4. Abu Bakar A.E. Determination of heavy metals (cadmium, copper, lead and nickel) in slimming teas by using atomic absorption spectroscopy. (2008);
5. Reena S, Neetu G, Anurag M and Rajiv G. Heavy metals and living systems An overview. Indian J Pharmacol. (2011); 43(3): 246-253.
6. Burkitt H G, Young B and Health J W. In Wheater's Functional Histology. London: Churchill Livingstone (2000); (4):282-6.
7. McCane K L and Huehe S.E. Pathophysiology: The biologic basis for disease in adults and children. (1994); New York (2)1320-1401.
8. Chugh K.S, Singhal P.C and Sharma, B.K. Methemoglobinemia in acute copper sulphate poisoning. (1975); Annals of Internal Medicine 82: 226.
9. Druet P, Bernard A. and Hersch F. et al. Immunologically mediated glomerulonephritis induced by heavy metals. Archives of Toxicology. (1982); 50 (3-4):187-194.
10. Fuentealba I and Haywood S. Cellular mechanisms of toxicity and tolerance in the copper-loaded rat. I. Ultra structural changes in the liver. (1988); Liver 8(6): 372-380.
11. Lei R, Wu C, Yang B, Ma H, Shi C, Wang Q, Yuan Y and Liao M. Integrated metabolomic analysis of the nano-sized copper particle-induced hepatotoxicity and nephrotoxicity in rats. A rapid in vivo screening method for nanotoxicity. Toxicology and Applied Pharmacology (2008); 232 (2): 292-301.
12. Bertinato J and Zouzoulas A. Considerations in the development of biomarkers of copper status. (2009); J.AOAC Int. 92 (5): 1541-1550.
13. Onwuka Gl. Food Analysis and Instrumentation. Theory and Practice. Napthali Prints. (2005). p. 140-146.
14. Freireich E J, Gehan EA, Rall D P, Schmidt LH and Skipper H . E. Quantitative comparison of toxicity of anticancer agents in mouse, rat, dog, monkey and man. Cancer Chemotherap Rep., (1966); 50(4):219-244.
15. Cavaggioni A, Mucignat-Caretta C. "Major urinary proteins, alpha (2U)-globulins and aphrodisin". Biochimica et Biophysica Acta (2000); 1482 (1-2): 218-28. doi:10.1016/S0167-4838(00)00149-7. PMID 11058763
16. Cunliffe-Beamer, T. L. Biomethodology and surgical techniques. (1983); Pp. 419-420 in The Mouse in Biomedical Research, Vol III, Normative Biology, Immunology and Husbandry. H. L. Foster, J. D. Small and J. G. Fox, eds. New York: Academic Press
17. Sinkovic A, Strdin A and Svenssek F. Severe acute copper sulphate poisoning: A case report. Arh Hig Rada Toksikol. 2008; 59(1): 31-5.
18. Galhardi CM, Diniz YS, Faine LA, et al. Toxicity of copper intake: Lipid profile, oxidative stress, and susceptibility to renal dysfunction. Food Chem Toxicol. (2004); 42(12): 2053-60.
19. Wang Y, Katzmarzyk PT, Horswell R, Zhao W, Johnson J and Hu G. Kidney function and the risk of cardiovascular disease in patients with type 2 diabetes. Kidney International (2014); 85, 1192-1199.
20. Babaknejad and N Moshtaghie A.A. and Shahanipour K. The Toxicity of Copper on Serum Parameters Related to Renal Functions in Male Wistar Rats. (2015); Journal of Research in Medical Sciences.
21. Beckett JM, Hartley TF and Ball MJ. Evaluation of the Randox colorimetric serum copper and zinc assays against atomic absorption spectroscopy. Annals of Clinical Biochemistry. (2009); 46 (4):322-326.