

Association of High Cholesterol and Triglyceride Level in patients with Diabetes, Hypertension and Cerebrovascular Accidents

NAZIR AHMAD¹, MUHAMMAD NAVEED ASLAM², MUJTABA JAFFARY³

ABSTRACT

Aim: To assess the prevalence of high cholesterol and triglycerides in patients with diabetes mellitus, hypertension and cerebrovascular accident.

Methods: This study was carried out in the District Teaching Hospital, Ghazi Khan Medical College, DG Khan from June 2014 to May 2015. A total of 150 cases were included in the study.

Results: Out of 150 patients 93(62%) showed isolated stroke, 57(38%) showed isolated diabetes mellitus. Isolated hypertension was found in 21(7%), 6(2%) patients showed combined hypertension, CVA and diabetes mellitus. 18(6%) patients exhibited combined hypertension and stroke. 20(6.7%) patients were having combined diabetes mellitus and hypertension and 7(2.3%) patients exhibited combined diabetes mellitus and stroke.

Keywords: Diabetes mellitus, Hypertension, Cerebrovascular accident

INTRODUCTION

Cholesterol is a fatty material, which over the years thickens and narrows the arterioles. It is manufactured in the liver and is an essential element of blood. It is present abundantly in egg yolk, butter, cream, animal fat and milk. Hypertensive retinopathy is among the vascular complications of essential hypertension. It is known that the auto-regulation of retinal circulation fails as blood pressure increases beyond a critical limit. However, elevated blood pressure alone does not fully account for the extent of retinopathy¹.

There are various risk factors such as age, gender, familial trends, race and ethnic groups and modifiable factors such as hypertension, cardiac disease, diabetes mellitus, dyslipidaemia, smoking, alcohol abuse, physical inactivity, asymptomatic carotid stenosis and transient ischaemic attacks². Risk factors for strokes have been studied locally³. The relationship of serum lipids and lipoproteins with cerebrovascular disease are being studied along with many other risk factors as in coronary heart disease⁴. Several clinical trials showed an association between high concentrations of serum cholesterol and ischaemic stroke⁵.

The major risk factors contributing to the excess of cardiovascular disease caused by diabetes includes hyperglycemia, insulin resistance, dyslipidemia, hypertension and smoking.

Dyslipidemia in hypertensive patients is itself known to be a predisposing risk factor, an aggravating or complicating factor.⁶ Hypertension and hyperlipidemia not only accelerate atherogenesis but also cause degenerative changes in the walls of large and medium-sized arteries⁷, which accelerate cerebrovascular hemorrhage⁸, ischemic heart disease⁹, stroke and cardiac arrest¹⁰.

Increased serum cholesterol levels have been observed in individuals who suffer from non-haemorrhagic stroke, but not in those who suffer intra cranial or subarachnoid haemorrhage. Serum lipids are thought to interact with the pathogenesis of non-haemorrhagic stroke through a atherosclerosis mechanism.

PATIENTS AND METHODS

This study was carried out in the District Teaching Hospital, Ghazi Khan Medical College, DG Khan from June 2014 to May 2015. A total of 150 cases were included in the study.

RESULTS

Out of 150 patients, 76(50.7%) were male and 74(49.3%) were female. As regards dyslipidemia in CVA Hypertriglyceridemia was present in 18(46%) male and 21(54%) in female patients whereas hypercholesterolemia was seen in 9(60%) male and 6(40%) in female patients. Dyslipidemia in patients of hypertension, hypertriglyceridemia was present in 12(67%) male and 6(33%) in female patients and hypercholesterolemia was seen in 6(67%) male and 3(33%) in female patients. Dyslipidemia in diabetic

¹Assistant Professor of Medicine, District Teaching Hospital, Ghazi Khan Medical College DG Khan,

²Asstt Prof. of Medicine, Multan Medical & Dental College Multan

³Asst Professor of Medicine, Continental Medical College, Lahore
Correspondence to Dr. Nazir Ahmad, e-mail: drmalik.nazir.ahmed@gmail.com

patients, hypertriglyceridemia was present in 15(38%) male and 24(68%) in female patients, hypercholesterolemia was seen in 12 (50%) male and 12 (50%) in female patients.

Table 1: Triglycerides and cholesterol level in CVA (n=300 with CVA 48)

Age (yrs)	Male	Female
< 40	05	05
41-50	04	04
51-60	13	07
61-70	03	05
>70	02	-
Total	27	21

Table 2: Triglycerides and cholesterol level in patients with hypertension (n=300 with hypertension 23)

Age (yrs)	Male	Female
< 40	04	02
41-50	-	-
51-60	05	03
61-70	02	05
>70	02	-
Total	13	10

Table 3: Triglycerides and cholesterol level in patients with diabetes (n=300 with diabetes 60)

Age (yrs)	Male	Female
< 40	18	06
41-50	05	03
51-60	14	08
61-70	04	02
>70	-	-
Total	41	19

DISCUSSION

Stroke is a clinical syndrome characterized by rapidly developing symptoms and/or signs of focal and at times global loss of cerebral functions, with symptoms lasting more than 24 hours or leading to death with no apparent cause other than that of vascular origin.¹¹ According to World Health Organization report for 2002, total mortalities due to stroke in Pakistan were 78512.¹²

In the present study 60 (36.4%) patients of the total 150 patients with mean age of 54 years were suffering from isolated diabetes mellitus. Of these 60 patients, 45% were male while 55% were female. In this study 21% of the patients were having increased level of serum cholesterol and 34.2% patients having increased level of triglycerides (>150 mg/dl). When we compare this study with a study, it was found that 24% of the population was smoker and 64% of the patients were having sedentary life style.¹³

Another study showed that the effect of flavastatin HMG CoA reductase inhibitor was assessed in 50 patients with dyslipidemia in diabetic

belonged to the city of Lahore.¹⁴ In this study the total cholesterol and triglyceride were high. When compare this study with the present one, total cholesterol and triglycerides are quite low in present study. The reason is financial status, difference between people belonging to Lahore and our area.

Hyperglyperidemia is most common lipid abnormality in NIDDM, particularly in diabetics with poor glycemic control¹⁵. The above observation was also seen in present study. Triglyceride levels were above the desired level in 34.2% patients out of 60 patients, with mean triglyceride level below 150 mg/dl. Hyperclyceridemia is a potent risk factor for macrovascular disease due to reduced synthesis of insulin dependent lipoprotein lipase in liver, resulting in impaired clearance of LDL, cholylicron and VLDL remnants¹⁶. High triglyceride levels are associated with hypercoagulopathy and decreased fibrinolysis both contributing to coronary heart disease¹⁷. In addition to metabolic relation to LDL cholesterol modifying its particle into more dense, small and hence more atherogenic form¹⁸.

In the present study 23 (15.3%) patients with mean age of 63 years were suffering from isolated hypertension. Of these 57% were male while 43% had increased level of triglyceride. In these patients 32% were smoker. The similar study showed that 16% of the patients were having hyperglyceridemia and 5% of patients were having both raised triglyceride and cholesterol¹⁹.

In our study more patients had increased level of cholesterol and triglyceride. The exact cause of increased cholesterol and triglyceride in present study is not known. A study carried out reveals that 48% of patients were having hypercholesterolemia²⁰. Another study showed that 50% of the patients had dyslipidemia and 39% had hypertriglyceridemia²¹.

REFERENCES

1. Liebreich R. Bright's disease at diagnosis-shear ophthalmoscopy Albercht Von Graefes. Arch Ophthalmol 2007; 859: 5:265-8.
2. Khan J, Attique-ur-Rehman, Ali SA, Jielani A. Frequency of hypertension in stroke patients presenting at Ayub Teaching Hospital. J Ayub Med Coll Abbottabad 2006; 18:59-61.
3. Iqbal F, Hussain S, Hassan M. Hypertension, diabetes mellitus and hypercholesterolaemia as risk factors for stroke. Pak J Med Res 2003; 42:17-22.
4. Bashir K, Langhorne P, Lees KR, MacAlpine C, Muir K, Murray S, et al. Epidemiological aspects of referral to TIA clinics in Glasgow. Scott Med J 2007; 52:4-8.
5. Ali L, Jameel H, Shah MA. Risk factors in stroke. J Coll Physicians Surg Pak 1997; 7:7-10.
6. Thakur AK, Achari V. A study of lipid levels in uncomplicated hypertension. Indian Heart J 2000; 52: 173-7.

7. Naomi DL, Gordon HW. Hypertensive Vascular Disease. In: Fisher ND, Williams GH, editors. *Harrison's Principles of Internal Medicine*. 16th Ed. New York: McGraw Hill; 2005. p. 1463-70
8. Marshal SM, Flvvbieryg A. Prevention and early detection of vascular complications of diabetes. *BMJ* 2006; 333(7566): 475-80.
9. Buse JB, Bigger JT, Bvington RP. Action to control cardiovascular risk in diabetes (ACCORD) trial: design and methods. *Am J Cardiol* 2007; 99(12):21-33.
10. Ye J, Kiage JN, Arnett DK, Bartolucci AA, Kabagambe EK. Short-term effect of fenofibrate on C-reactive protein. *Diabetol Metab Syndr* 2011; 3: 24.
11. Kurozumi A, Okada Y, Mori H, Arao T, Tanaka Y. Efficacy of α -glucosidase inhibitors combined with dipeptidyl-peptidase-4 inhibitor. *J Diabetes Investig* 2013; 4(4): 393-8.
12. Ferreira M, António N, Gonçalves F. Hemoglobin: simply a laboratory value or a powerful predictor of risk in patients with acute coronary syndrome?. *Rev Port Cardiol* 2012; 31(2): 121-31.
13. Duncan BB, Wong TY, Tyroler HA, Davis CE, Fuchs FD. Hypertensive retinopathy and incident of coronary heart disease in high risk men. *BR J Ophthalmol* 2002; 86: 1002-6.
14. Ye J, Kiage JN, Arnett DK, Bartolucci AA, Kabagambe EK. Short term effect of fenofibrate on C-reactive protein. *Diabetol Metab Syndr* 2011; 3: 24.
15. Kuwashiro T, Sugimori H, Ago T, Kamouchi M, Kitazono T. Risk factors predisposing to stroke recurrence within one year of non-cardioembolic stroke onset. *Cerebrovasc Dis* 2012; 33(2): 141-9.
16. Ferreira M, Antonio N, Goncalves F. Hemoglobin: simply a laboratory value or a powerful predictor of risk in patients with acute coronary syndrome. *Rev Port Cardiol* 2012; 31(2):121-31.
17. Hui ML, Kumar A, Adams GG. Protocol-directed insulin infusion sliding scales improve perioperative hyperglycaemia in critical care. *Perioper Med (Lond)* 2012; 1: 7.
18. Kuwashiro T, Sugimori H, Ago T, Kamouchi M, Kitazono T. Risk factors predisposing to stroke recurrence within one year of non-cardioembolic stroke onset. *Cerebrovasc Dis* 2012; 33 (2): 141-9.
19. Ikeda N, Iijima R, Hara H, Moroi M, Nakamura M, Sugi K. Glycated hemoglobin is associated with the complexity of coronary artery disease, even in non-diabetic adults. *J Atheroscler Thromb* 2012; 19(12): 1066-72.
20. Stenestrand U, Wijkman M, Fredrikson M, Nystrom FH. Association between admission supine systolic blood pressure and 1-year mortality. *JAMA* 2010; 303(12): 1167-72.
21. Sandercock P, Mielke O, Liu M, Counsell C. Anticoagulants for preventing recurrence following presumed non-cardioembolic ischaemic stroke or transient ischaemic attack. *Cochrane Database Syst Rev* 2003; (1): CD000248