

Prevalence of Hyperkalemia in Myocardial infarction

MEMOONA MUMTAZ¹, MIR AHMED RAZA², IMRAN ALI³

ABSTRACT

Aim: Increased potassium level in cardiac patient is extremely dangerous. It should be controlled strictly in myocardial infarction patients with precautions.

Methods: Total 3340 patients with hyperkalaemia history were collected from Mayo hospital Lahore. Majority of patients are hospital admitted. Mean age 49±5 years, 59% males and 41% females. We assess the incidence and influencing factors on potassium level in myocardial infarction patients. Patients were classified as : <4 mEq/L, 4 to <4.8 mEq/L, 4.8 to <5.5, 5.5 to <6.0 mEq/L, and ≥6.0 mEq/L. Linear regression analysis were applied on multivariate variables. The association between the increased risk of hyperkalemia in hospital admitted patients and mortality rate were also assessed.

Results: From 3340 patients with hyperkalemia with or without myocardial infarction. The hyperkalemia rate (potassium <4.8 mEq/L) was 19.3% in patients with chronic cardiac diseases. Mild to moderate hyperkalemia potassium rate (potassium 5.5 to <6.0 mEq/L) had in 11.3% of patients. The most high rate of hyperkalemia (potassium ≥ 6.0 mEq/L) in patients was in 23%. There were unreasonable high mortality risk were noted in high rate of hyperkalemia in patients. There is strong relationship with early diagnosed patients and excellent prognosis.

Conclusion: Hyperkalemia is very dangerous in cardiac disease patients. Higher levels of potassium in patients are directly coupled with increased mortality rate in hospitals, with increased risks of hyperkalemia adverse outcomes were noted even at mild rate of hyperkalemia. It is concluded that more aggressive and quick treatment of hyperkalemia may improve patient health and prognosis.

Keywords: Hyperkalemia, myocardial infarction, Cardiac disease.

INTRODUCTION

Increase rate of potassium in human body can progress to grim and vitally fatal cardiac rhythm irregularities. Earlier research data has strongly suggested the influence of hyperkalemia on dangerous cardiac results¹. With the advancement in medical sciences, it is becomes prominent that incidence of chronic kidney disease and diabetes both increases in hyperkalemic patients². Moreover, most of the acute myocardial infarction cases has increased hyperkalemia with or without happening of heart failure³.

The high rate of using beta-blockers and rennin angiotensin aldosterone system antagonists has been associated with increase rate of hyperkalemia in patients and become major cause of hospitalization and deaths⁴. Moreover, there is many surgical intervention like coronary artery bypass grafting and percutaneous procedures of myocardial infarction may circuitously leading to risk of hyperkalemia through damaging of nephropathy and acute kidney injury⁵.

On the other hand earlier studies have measure the direct relationship between the potassium level in

blood and rate of mortality following acute myocardial infarction, no one describe the existing prevalence and prognostic value of many factors of hyperkalemia in patients⁶. Furthermore, it is totally undefined the role of hyperkalemia results in enhancing the patient outcomes⁷. Additional describing the risk linked with hyperkalemia on all body health, and with associated other diseases, may lead to improve patients health with many benefits⁸.

PATIENTS & METHODS

Total 3340 patients with high serum potassium history were collected from Mayo hospital Lahore. Patients hospitalized between June 2015 to June 2017. Majority of patients are hospital admitted. Data were sort out according to demographic characteristics, complete medical history, laboratory tests, surgical procedures, medication and complications. All the documentation of hyperkalemic patients were categorize after diagnosis by the cardiac physicians. All the patients who had whole blood potassium increases with the serum potassium values for this study. Patients barred from participation in study were those admitted from hospice or with history of trauma.

Mean age 49±5 years, 59% males and 41% females. We assess the incidence and influencing factors on potassium level in myocardial infarction

¹MO Accident & Emergency, Sheikh Zayed Hospital, RYK.

²MO BHU Harria Tehsil Malikwal Dist. Mandibahauddin

³MO BHU Chak 16 Tehsil Malakwal Dist. Mandibahauddin

Correspondence to Dr. Memoona Mumtaz E-mail: memoonaosama.pansota@gmail.com

patients. Patients were classified as : <4 mEq/L, 4 to <4.8 mEq/L, 4.8 to <5.5, 5.5 to <6.0 mEq/L, and ≥6.0 mEq/L. Linear regression analysis were applied on multivariate variables. The association between the increased risk of hyperkalemia in hospital admitted patients and mortality rate were also assessed. The association between higher K and high mortality risk were persisted with multivariate variables. In addition patients with high hyperkalemia values experienced high rate in hospital admission.

RESULTS

From 3340 patients with hyperkalemia with or without myocardial infarction. The hyperkalemia rate (potassium <4.8 mEq/L) was 19.3% in patients with chronic cardiac diseases. Mild to moderate hyperkalemia potassium rate (potassium 5.5 to <6.0 mEq/L) had in 11.3% of patients. The most high rate of hyperkalemia (potassium ≥ 6.0 mEq/L) in patients was in 23%. Continuity of variables were compared using a straight trend test while categorical variables were measured using mantel trend test. There were unreasonable high mortality risk were noted in high rate of hyperkalemia in patients. There is strong relationship with early diagnosed patients and excellent prognosis. In addition, patients with late diagnosis and higher rate of hyperkalemia shows high mortality rate in mayo hospital lahore. The high number of patients with maximum potassium per patients were classified according to their complication and shows much variations in concluding of results.

Table 1: Hospital mortality rate by number of hyperkalemia values.

Hyperkalemia Values	Mortality Rate	P-Value
0 value ≥ 5	3.2%	<0.01
1 value ≥ 5	12.6%	<0.01
2 values of ≥ 5	15.4%	<0.01
3 values of ≥ 5	21.9%	<0.01

DISCUSSION

In this study, we analyze that hyperkalemia found in estimate more than twenty percent of patients on dialysis and more than 66% were found in non-dialysis patients. There were high mortality rate observed with higher potassium level, specifically in patients with secondary diseases. Moreover, patients admitted in hospital was more in percentage with high potassium than any other cardiac issue(9). To our information, this study is among the very first to describe the prevalence of hyperkalemia, the increasing rate of inpatients mortality linked with other cardiovascular diseases(10).

In past studies, results of the hyperkalemia in cardiovascular diseases have much most variations. Goyal et al, examined the relationship of potassium levels with hospital admitted patients with ventricular arrhythmias, cardiac arrest and searching an increase risk of hospital admitted patients among high potassium level patents¹¹. An older previous study including 956 patients with myocardial infarction and presented relationship between hyperkalemia patients and early post infarction ventricular events¹².

Many researchers have concluded increased hyperkalemia events coupled with common adaption of medications including beta blockers, angiotensin converting enzyme and aldosterone receptors blockers¹³. Our study describe the earlier precautions to decrease the mortality rate with association of varying degrees of potassium in patients¹⁴. While treatment values are not evaluated in this study, but it is significant that diagnosis of hyperkalemia within time and its preventive measure give better opportunity to improve prognosis then persistent hyperkalemia¹⁵.

We noticed that late or poor diagnosis linked with hyperkalemia may give clinicians tough time in evaluation of patients for modifiable factors that are linked with hyperkalemia and following any secondary diseases¹⁶. In the methods of management point of view. Several new products for management are in development, which may provide proper intime treatment of patients with better outcomes¹⁷.

Although these novel treatment modalities may ultimately change the way hyperkalemia is managed in the hospital, they are still undergoing investigation in the acute and outpatient settings¹⁸. Our study is particularly timely in light of development of new medications for the management of hyperkalemia, and highlights an opportunity to investigate the heterogeneity of treatment benefit of hyperkalemia patients¹⁹.

CONCLUSION

Hyperkalemia is very dangerous in cardiac disease patients. Higher levels of potassium in patients are directly coupled with increased mortality rate in hospitals, with increased risks of hyperkalemia adverse outcomes were noted even at mild rate of hyperkalemia. It is concluded that more aggressive and quick treatment of hyperkalemia may improve patient health and prognosis.

REFERENCES

1. An JN, Lee JP, Jeon HJ, Kim DH, Oh YK, Kim YS, et al. Severe hyperkalemia requiring hospitalization: predictors of

- mortality. *Critical care* (London, England). 2012 Nov 21;16(6):R225. PubMed PMID: 23171442. Pubmed Central PMCID: PMC3672605. Epub 2012/11/23. eng.
2. McCullough PA, Beaver TM, Bennett-Guerrero E, Emmett M, Fonarow GC, Goyal A, et al. Acute and chronic cardiovascular effects of hyperkalemia: new insights into prevention and clinical management. *Reviews in cardiovascular medicine*. 2014;15(1):11-23. PubMed PMID: 24762462. Epub 2014/04/26. eng.
 3. Weir MA, Juurlink DN, Gomes T, Mamdani M, Hackam DG, Jain AK, et al. Beta-blockers, trimethoprim-sulfamethoxazole, and the risk of hyperkalemia requiring hospitalization in the elderly: a nested case-control study. *Clinical journal of the American Society of Nephrology : CJASN*. 2010 Sep;5(9):1544-51. PubMed PMID: 20595693. Pubmed Central PMCID: PMC2974392. Epub 2010/07/03. eng.
 4. Pitt B, Bakris G, Ruilope LM, DiCarlo L, Mukherjee R. Serum potassium and clinical outcomes in the Eplerenone Post-Acute Myocardial Infarction Heart Failure Efficacy and Survival Study (EPHESUS). *Circulation*. 2008 Oct 14;118(16):1643-50. PubMed PMID: 18824643. Epub 2008/10/01. eng.
 5. Juurlink DN, Mamdani MM, Lee DS, Kopp A, Austin PC, Laupacis A, et al. Rates of hyperkalemia after publication of the Randomized Aldactone Evaluation Study. *The New England journal of medicine*. 2004 Aug 05;351(6):543-51. PubMed PMID: 15295047. Epub 2004/08/06. eng.
 6. Wozakowska-Kaplon B, Janowska-Molenda I. Iatrogenic hyperkalemia as a serious problem in therapy of cardiovascular diseases in elderly patients. *Polskie Archiwum Medycyny Wewnetrznej*. 2009 Mar;119(3):141-7. PubMed PMID: 19514643. Epub 2009/06/12. eng.
 7. Perazella MA. Drug-induced hyperkalemia: old culprits and new offenders. *The American journal of medicine*. 2000 Sep;109(4):307-14. PubMed PMID: 10996582. Epub 2000/09/21. eng.
 8. Fordjour KN, Walton T, Doran JJ. Management of hyperkalemia in hospitalized patients. *The American journal of the medical sciences*. 2014 Feb;347(2):93-100. PubMed PMID: 23255245. Epub 2012/12/21. eng.
 9. Salisbury AC, Reid KJ, Marso SP, Amin AP, Alexander KP, Wang TY, et al. Blood transfusion during acute myocardial infarction: association with mortality and variability across hospitals. *Journal of the American College of Cardiology*. 2014 Aug 26;64(8):811-9. PubMed PMID: 25145526. Epub 2014/08/26. eng.
 10. McMahon GM, Mendu ML, Gibbons FK, Christopher KB. Association between hyperkalemia at critical care initiation and mortality. *Intensive care medicine*. 2012 Nov;38(11):1834-42. PubMed PMID: 22806439. Epub 2012/07/19. eng.
 11. Goyal A, Spertus JA, Gosch K, Venkitachalam L, Jones PG, Van den Berghe G, et al. Serum potassium levels and mortality in acute myocardial infarction. *Jama*. 2012 Jan 11;307(2):157-64. PubMed PMID: 22235086. Epub 2012/01/12. eng.
 12. Weir MR, Bakris GL, Bushinsky DA, Mayo MR, Garza D, Stasis Y, et al. Patiromer in patients with kidney disease and hyperkalemia receiving RAAS inhibitors. *The New England journal of medicine*. 2015 Jan 15;372(3):211-21. PubMed PMID: 25415805. Epub 2014/11/22. eng.
 13. Kosiborod M, Rasmussen HS, Lavin P, Qunibi WY, Spinowitz B, Packham D, et al. Effect of sodium zirconium cyclosilicate on potassium lowering for 28 days among outpatients with hyperkalemia: the HARMONIZE randomized clinical trial. *Jama*. 2014 Dec 03;312(21):2223-33. PubMed PMID: 25402495. Epub 2014/11/18. eng.
 14. Packham DK, Rasmussen HS, Lavin PT, El-Shahawy MA, Roger SD, Block G, et al. Sodium zirconium cyclosilicate in hyperkalemia. *The New England journal of medicine*. 2015 Jan 15;372(3):222-31. PubMed PMID: 25415807. Epub 2014/11/22. eng.
 15. Madias JE, Shah B, Chintalapally G, Chalavarya G, Madias NE. Admission serum potassium in patients with acute myocardial infarction: its correlates and value as a determinant of in-hospital outcome. *Chest*. 2000 Oct;118(4):904-13. PubMed PMID: 11035655. Epub 2000/10/18. eng.
 16. Nordrehaug JE, Johannessen KA, von der Lippe G. Serum potassium concentration as a risk factor of ventricular arrhythmias early in acute myocardial infarction. *Circulation*. 1985 Apr;71(4):645-9. PubMed PMID: 3971535. Epub 1985/04/01. eng.
 17. Macdonald JE, Struthers AD. What is the optimal serum potassium level in cardiovascular patients? *Journal of the American College of Cardiology*. 2004 Jan 21;43(2):155-61. PubMed PMID: 14736430. Epub 2004/01/23. eng.
 18. Kosiborod M, Inzucchi SE, Goyal A, Krumholz HM, Masoudi FA, Xiao L, et al. Relationship between spontaneous and iatrogenic hypoglycemia and mortality in patients hospitalized with acute myocardial infarction. *Jama*. 2009 Apr 15;301(15):1556-64. PubMed PMID: 19366775. Epub 2009/04/16. eng.
 19. Hofsten DE, Wachtell K, Lund B, Molgaard H, Egstrup K. Prevalence and prognostic implications of non-sustained ventricular tachycardia in ST-segment elevation myocardial infarction after revascularization with either fibrinolysis or primary angioplasty. *European heart journal*. 2007 Feb;28(4):407-14. PubMed PMID: 17227787. Epub 2007/01/18.