

Examine the Prevalence of Oral Squamous Cell Carcinoma also Determine the Risk Factors and Causes of Improper Diagnosis

RUMMANA AQEEL¹, ZEESHAN ASLAM², ANEELA AMJAD³, AMMARA ANWAR⁴, FAISAL ASGHAR⁵, TAHMASUB FARAZ TAYYAB⁶

¹Senior Registrar Oral Pathology Avicenna Medical & Dental College Lahore,

²Senior Registrar Oral Medicine, Rashid Latif Dental College Lahore,

³Assistant Professor of Oral Medicine & Diagnostic Sharif Medical & Dental College Lahore

⁴Senior Registrar Oral Medicine Rashid Latif Dental College Lahore

⁵Assistant Professor of Oral & Maxillofacial Surgery, Multan Medical & Dental College Multan

⁶Assistant Professor of Oral & Maxillofacial Surgery UCMD, University of Lahore

Correspondence to Dr. RummanaAqeel

ABSTRACT

Aim: To examine the prevalence of clinical and histological proven cases of oral-squamous cell carcinoma also examine the risk factors and challenges in diagnosis oral cancers.

Study Design: Retrospective/observational

Place & duration: Department of Oral Pathology, Ghurki Teaching Hospital Lahore from 1st April 2016 to 31st January 2017.

Methods: Total sixty patients of both genders with ages 30 to 70 years clinical and histologically diagnosed to have OSCC (oral squamous cell carcinoma) were included in this study. Patients demographical details including age sex, residence, socio-economic status were examined after taking informed consent. Incisional biopsy was taken from all the patients and sent to laboratory for examination. Prevalence and risk factors of OSCC were examined. Challenges in diagnosing were also examined.

Results: Out of 60 patients 45 patients found to have OSCC in which 35 (77.76%) were males while 10(22.22%) were females. 20(44.44%) patients were ages 30 to 45 years, 19(42.22%) patients had ages 46 to 60 years and 6(13.33%) patients were ages above 60 years. Buccal mucosa was the most frequent site of OSCC found in 23(51.11%) patients followed by lower alveolar and tongue. The most frequent risk factor was cigarette smoking found in 18(40%) patients. delay due to patients unawareness found in 15(33.33%) patients followed by misdiagnosed by expertise and lack of facility (Diagnosing tools).

Conclusion: There is a high prevalence of oral cancer and smoking is the major risk factor of this malignant disease. Misdiagnosed at first visit is the major concern, Lack of diagnosis facilities and lack of awareness are also most important factor for increasing the rate of this malignant disorder.

Keywords: Oral Squamous cell carcinoma (OSCC), Frequency, Risk factors, Diagnosis, Causes

INTRODUCTION

Globally, Oral cancer is considered most frequent life threatening malignant disorder and from last thirty years the incidence rate of oral cancer is going upward. Due to oral cancer the high rate of mortality is recorded and in Pakistan oral cancer contributed a great threat to public health with high rate of mortality and morbidity¹. As per international reports oral cancer is the most common malignant disorder and in Pakistan it is the 2nd most common malignant disease that lead to increase rate of mortality.² From all the types of tumor, oral cancer in Pakistan rated 15% and this rate is 5 times greater than the worldwide incidence rate of oral cancer 3%³. According to the international researches, In sub-continent oral cancer incidence rate is quite high with incidence rate of 7.9% and mortality rate 3.8%⁴.

Many of studies reported that the patients with third to 5th decade of life has a high prevalence of oral cancer but now a days several studies reported early age. It is reported that majority of oral cancers are squamous cell carcinoma and accounted above 90%³⁻⁴.

Many of factors involves in raising the incidence of oral squamous cell carcinoma including genetic and environmental factors. Tobacco use, betel quid, alcohol, chewing tobacco and snuff are the most frequent risk factors⁵.

Worldwide, tobacco use and alcohol consumption considered the strongest risk factors and in developed countries these two factors are most common and contributed a high rate of incidence.^{6,7} In developing countries like Pakistan, India, Bangladesh tobacco use with betel quid and snuff with substitute are the major risk factors of oral squamous cell carcinoma.⁸ Globally, early and accurate diagnosis of oral cancer is a big challenge for professional. Presenting instances in which pathologies were not diagnosed and/or treated in spite of obvious signs, should serve as a warning for dental professionals. The general practitioner is the "first line of defense" for symptomatic and asymptomatic patients and therefore the implementation of well-established screening protocols is of paramount importance. In the light of the low sensitivity ratio of oral examinations, use of biopsies is mandatory when a lesion is suspected.⁹⁻¹²

Received on 13-04-2019

Accepted on 23-07-2019

MATERIALS AND METHODS

This retrospective analysis was conducted at Department of Oral Pathology Ghurki Teaching Hospital Lahore and the duration of study was 10 months, from April 2016 to Jan 2017. In this study total sixty patients of both genders with ages 30 to 70 years clinical and histologically diagnosed to have oral squamous cell carcinoma were included in this study. Patients demographical details including age sex, residence, socio-economic status were examined after taking informed consent. Patients with other severe infections, patients diagnosed to had no oral cancer and those who were not interested to participate were excluded from this study. All the patients had undergone incisional biopsy. A tissue were obtained and sent to laboratory for histopathological examination of the oral squamous cell carcinoma. Microscopic examination was done to examine the stages of tumor. Risk factors were also examined. Causes of severe malignancy were recorded. All the data was analyzed by SPSS 20.

RESULTS

Out of 60 clinically proven cases 45 patients found to have squamous cell carcinoma by histopathological examination. In which 35(77.78%) were males while 10(22.22%) were females. 20(44.44%) patients were ages 30 to 45 years, 19 (42.22%) patients had ages 46 to 60 years and 6(13.33%) patients were ages above 60 years. 25 (55.56%) patients had rural residency while 20(44.44%) had urban residency. 23(51.11%) patients had low socio-economic status, 22(48.89%) had middle status (Table 1). According to pathological examination, 23(51.11%) patients had buccalmucosa, 12(26.67%) patients had lower alveolar mucosa, 4(8.89%) patients had tongue, 3(6.67%) had lips and 3(6.67%) patients had cheeks according to the sites of tumor (Table 2).

The most frequent risk factor was cigarette smoking found in 18(40%) patients followed by betel quid 12(26.67%), alcohol 10(22.22%), 3(6.67%) smoking with tobacco chewing and 2(4.44%) had others (Table 3). Most important cause of malignancy was misdiagnosed by expertise at first visit found in 13(28.89%), delay due to patients unawareness found in 15(33.33%), delay in diagnosis due to lack of facility (Diagnosing Tools found in 6(13.33%) (Table 4).

Table 1: Demographical characteristics of all the patients

Characteristics	No.	%
Gender		
Male	35	77.78
Female	10	22.22
Age (years)		
30 – 45	20	44.44
46 – 60	19	42.23
> 60	6	13.33
Residence		
Urban	20	44.44
Rural	25	55.56
Socioeconomic status		
Low	23	51.11
Middle	22	48.89

Table 2: Sites of oral squamous cell carcinoma

Sites	No.	%
Buccal Mucosa	23	51.11
Alveolar	12	26.67
Tongue	4	8.88
Cheeks	3	6.67
Lips	3	6.67

Table 3: Risk factors of oral squamous cell carcinoma

Risk Factors	No.	%
Cigarette smoking	18	40
Betel Quid	12	26.67
Alcohol	10	22.22
Smoking with Tobacco chewing	3	6.67
Other	2	4.44

Table 4: Causes of malignant oral tumor associated to diagnosing

Causes	No.	%
Misdiagnosed by expertise	13	28.89
Patients Unawareness	15	33.33
Lack of Diagnosing Tools	6	13.33
Others (faculty, procedure)	3	6.67

DISCUSSION

OSCC is found to be more common in slightly older males all over the world but newer studies are revealing its development in younger age groups.¹³ Many of studies reported that the patients with third to 5th decade of life has a high prevalence of oral cancer but in recent years several studies reported early age. Some previous studies shows that patients with age group 50 to 60 years had a high prevalence of OSCC^{14,15,16}. The present study was conducted aimed to examine the frequency and risk factors of oral squamous cell carcinoma also determine the causes of this malignant disorder associated to diagnosis. In our study majority of patients were males 77.78% while female patients were very low in numbers. These results showed similarity to some other studies in which male patients population was high 60 to 85% as compared to females^{17,18}.

In present study we found that majority of patients were ages between 30 to 60 years. A study conducted by Sadia et al¹⁹ regarding oral squamous cell carcinoma reported patients with ages 30 to 60 years had a high prevalence of oral tumor. We found that 25(55.56%) patients had rural residency while 20(44.44%) had urban residency. 23(51.11%) patients had low socio-economic status, 22(48.89%) had middle status. These results were comparable to some other studies in which most of the patients had low socio-economic status. As per previous studies socio-economic status and literacy contributed a lot for increasing the rate of oral tumors²⁰⁻²². In this study, we found as per pathological examination 23(51.11%) patients had buccal mucosa, 12(26.67%) patients had lower alveolar mucosa. Most of the studies demonstrated buccal mucosa was the most frequent site of oral squamous cell carcinoma followed by alveolar mucosa²³⁻²⁴.

We found that smoking tobacco (Cigarettes) was the most frequent risk factor for developing oral cancer 40% followed by betel quid and alcohol. These results were similar to some other studies but in contrast many of studies reported alcohol consumption, snuff and chewing tobacco were the most common risk factors.²⁵⁻²⁶ The second most important objective of this study was to

examine the causes, challenges associated to diagnosing oral cancers and we found most important cause of malignancy was misdiagnosed by expertise at first visit found in 13(28.89%), delay due to patients unawareness found in 15(33.33%), delay in diagnosis due to lack of facility (Diagnosing Tools found in 6(13.33%). Many of previous studies shows the problems regarding malignancy and found misdiagnosed by expertise and delay in diagnosis due to unawareness were the most important challenges for diagnosing this malignant disorder.²⁷⁻²⁸ In this study we observed that there is a need of trained professionals for diagnosing this malignant disorder also provide awareness to the people so that early diagnosis could be possible and may helps to reduce the morbidity and mortality rate in Pakistan.

CONCLUSION

Oral cancers are the words most common and life threatening malignant disorders. In Pakistan the rate of oral cancers is quite high as compared to developed countries. In this study, we concluded that there is a high prevalence of oral cancer and smoking is the major risk factor of this malignant disease. Misdiagnosed at first visit is the major concern, Lack of diagnosis facilities and lacks of awareness are also most important factor for increasing the rate of this malignant disorder. Authorities should take a action for training the professionals and for providing the proper diagnosing tools so that mortality and morbidity could decrease. Also provide awareness to the people about this malignant disorder. Moreover, many of challenges contributed in raising the rate of oral cancer but importantly early diagnosis and prompt treatment are very helpful for decreasing the frequency of oral squamous cell carcinoma.

REFERENCES

1. Cancer Registry and Clinical Data Management (CRCDM) ShaukatKhanum Memorial Cancer Hospital and Research Center (SKMCH&RC) - (www.shaukatkhanum.org.pk). Report based on cancer cases registered at SKMCH&RC from Dec1994. Dec. 2011 and in 2011. Released June, 2012.
2. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010;127(12):2893–17.
3. Camargo CM, Voti L, Guerra M, Chapuis F, Mazuir M, Curado MP. Oral cavity cancer in developed and in developing countries: population-based incidence. *Head Neck* 2010;32(3):357–67.
4. Choi S, Myers JN. Molecular pathogenesis of oral squamous cell carcinoma: implications of therapy. *J Dent Res* 2008;87:14–32.
5. Chaturvedi AK. Epidemiology and clinical aspects of HPV in head and neck cancers. *Head Neck Pathol* 2012;6(1):S16–24.
6. Zygogianni AG, Kyrgias G, Karakitsos P. Oral Squamous cell cancer: early detection and the role of alcohol and smoking. *Head Neck Oncol* 2011;3
7. Saman DM. A review of the epidemiology of oral and pharyngeal carcinoma: update. *Head Neck Oncol* 2012;4
8. Lambert R, Sauvaget C, de Camargo Cancela M, Sankaranarayanan R. Epidemiology of cancer from the oral cavity and pharynx. *Eur J GastroenterolHepatol* 2011;23(8):633–641.
9. Warnakulasuriya S, Dietrich T, Bornstein MM, Peidro EC, Preshaw PM, Walter C. Oral health risks of tobacco use and effects of cessation. *Int Dent J* 2010;60(1):7–30.
10. Jerjes W, Upile T, Radhi H, Petri A, Abiola J, Adams A. The effect of tobacco and alcohol and their reduction/cessation on mortality in oral cancer patients: short communication. *Head Neck Oncol* 2012;4
11. Jayelekshmi P, Gangadharan P, Akiba S, Koriyama C, Nair RRK. Oral cavity cancer risk in relation to tobacco chewing and biri smoking among men in Karunagappally, Kerala, India: Karunagappally cohort study. *Cancer Sci* 2011;102(2):460–67.
12. Bhurgri Y, Bhurgari A, Nishta S, Ahmed A, Pervez S, Ahmed R, et al. Pakistan Country Profile of Cancer and Cancer Control 1995-2004. *J Pak Med Assoc* 2006;56(3):124–130.
13. Mafi N, Kadivar M, Hosseini N, Ahmadi S, Zare-Mirzaie A. Head and Neck Squamous Cell Carcinoma in Iranian patients and risk factors in young adults: a Fifteen-Year Study. *Asian Pacific J Prev* 2012;13(7):3373–3378.
14. Khan MA, Saleem S, Shahid SM, Hameed A, Qureshi NR, Abbasi Z, et al. Prevalence of oral squamous cell carcinoma (OSCC) in relation to different chewing habits in Karachi, Pakistan. *Pak J BiochemMolBiol* 2012;45(2):59–63.
15. Tanwir F, Altamash M, Gustafsson A. Influence of betel nut chewing, dental care habits and attitudes on perceived oral health among adult Pakistanis. *Oral Health Prev Dent* 2008;6(2):89-94.
16. Daud M et al. Oral squamous cell carcinoma (OSCC). *Pakistan Oral DentJ* 2016; 36(3): 13.
17. Dissanayaka WL, Pitiyage G, Kumarasiri PV, Liyanage RL, Dias KD, Tilakaratne WM. Clinical and histopathologic parameters in survival of oral squamous cell carcinoma. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2012;113:518–25.
18. Fukuda M, Ohmori Y, Sakashita H. The Role of Tumor Microenvironment in Oral Cancer. In: Biswas S, editor. *Tumor Microenvironment and Myelomonocytic Cells*. InTech 2012.
19. SadiaAkram et al. Emerging Patterns in Clinico-pathological spectrum of OralCancers. *Pak J Med Sci* 2013; 29(3): 783–787.
20. Rivera C, Venegas B. Histological and molecular aspects of oral squamous cell carcinoma (Review). *OncolLett* 2014;8:7–11.
21. van der Waal I. Potentially malignant disorders of the oral and oropharyngeal mucosa; terminology, classification and present concepts of management. *Oral Oncol* 2009;45:317–23.
22. Lee J, Taneja V, Vassallo R. Cigarette smoking and inflammation: cellular and molecular mechanisms. *J Dent Res* 2012;91:142–9.
23. Barnes L World Health Organization and Cancer IAFRo. Pathology and genetics of head and neck tumours. World Health Organization 2015.
24. National Cancer Institute Thesaurus. Available from: <http://ncit.nci.nih.gov/ncitbrowser/>. Accessed 13 June 2011.
25. Rojas-Alcayaga G, Brignardello R, Espinoza I, et al. Oral malignant tumours: a report on 728 cases in Chile. *ActaOdontolLatinoam* 2010;23:136–142.
26. Smith B, Goldberg LJ, Rutenber A, et al. Ontology and the future of dental research informatics. *J Am Dent Assoc* 2010;141:1173–75.