

Frequency of Spontaneous Bacterial Peritonitis in Liver Cirrhosis Patients Having Hepatic Encephalopathy

MUHAMMAD HAROON BILAL, MUHAMMAD TAHIR*, NAVEED ATIF KHAN**

ABSTRACT

Aim: To find out the frequency of SBP in patients with Hepatic Encephalopathy and association of different grades of hepatic encephalopathy with SBP.

Methods: This cross sectional study was conducted at Dera Ghazi Hospital, Dera Ghazi Khan from June 2013 to May 2014. Patients with hepatic encephalopathy and ascites with age from 20 to 60 years were included in this study. Patients were labeled as having SBP when ascetic fluid neutrophil count > 250/ml. Hepatic encephalopathy was graded according to west heaven criteria.

Results: Total 155 patients with hepatic encephalopathy were included in this study. Mean age of patients was 38.9±11.7. Males were 119(77%) and females were 36(23%). Overall SBP was present in 81(52.3%) patients.

Conclusion: HE and SBP are the common complications in liver cirrhosis. Most of the HE patients having SBP as well as precipitating factor. The treatment of SBP is the important step of management of HE in liver cirrhosis.

Keywords: Hepatic Encephalopathy, SBP

INTRODUCTION

Hepatic encephalopathy (HE) is a serious complication of advanced liver disease and refers to Neuropsychological syndrome that occurs in a patient with either severe liver failure, chronic liver disease or as a consequence of surgical or radiological portal systemic shunts.¹

Factors like gastrointestinal bleeding, infections, constipation, electrolyte imbalance (hyponatremia, hypokalemia), hypoglycemia, Medicines (sedative-hypnotics, opiates) could be responsible for the development of Hepatic encephalopathy. It is imperative to identify one of these factors so that prompt treatment given to get patient out of encephalopathy.^{2,3}

Ascites is defined as the collection of free fluid in the peritoneal cavity. More than 500ml of fluid must be present before ascites can be presented clinically.⁴

Spontaneous Bacterial Peritonitis(SBP) is defined as the infection of ascetic fluid without any apparent intra-abdominal focus of infection in patients of cirrhosis of liver.⁵ SBP is a common complication in patients of cirrhosis of liver⁶. The signs and symptoms of SBP are subtle compared with those of patients who have surgical peritonitis in the absence of ascites⁴. Ascites prevent the development of rigid

abdomen by separating the visceral pleura from the parietal pleura. Patients of cirrhosis of liver with SBP may be asymptomatic not having the clinical sign and symptoms of peritonitis like abdominal pain, abdominal discomfort and fever⁴. Despite the effective treatment of SBP, it has the high morbidity and mortality. Mortality rate of SBP is 70% per one year due to the complications like septic shock, hepatorenal syndrome and hepatic encephalopathy. Various international studies show that 10% to 30% of patients of cirrhosis of liver develop SBP⁷. One of the predisposing factors which are responsible for hepatic encephalopathy and subsequent deterioration in the condition of cirrhosis patient is appearance of spontaneous bacterial peritonitis (SBP)⁸.

SBP is characterized by abrupt onset of fever, chills abdominal pain with rebound tenderness over abdomen, absent bowel sounds and leucocytosis. Paracentesis reveals cloudy ascitic fluid with many WBCs predominantly, polymorphonuclear cells (PMN). SBP is an important cause of morbidity and mortality in patients with hepatic encephalopathy⁹. It is a serious and potentially life-threatening complication and associated with a 30–50% mortality rate, if not treated and death can occur within few hours¹⁰ so early diagnostic paracentesis should be performed in every patient presented with hepatic encephalopathy¹¹.

Keeping in view the high mortality rate of SBP in patients of liver cirrhosis, a study was conducted to find out the frequency of SBP in patients with hepatic

Assistant Professor Medicine, Dera Ghazi Khan Medical College, D.G. Khan

*Assistant Professor Pathology, Multan Medical & Dental College, Multan

**Assistant Professor Pathology, Dera Ghazi Khan Medical College, D.G. Khan

Correspondence to Dr. Muhammad Haroon Bilal, Assistant Professor E-mail: drmuhab@gmail.com

encephalopathy and to determine the association of different grades of hepatic encephalopathy with SBP.

MATERIAL AND METHODS

This cross sectional study was conducted at Dera Ghazi Hospital, Dera Ghazi Khan from June 2013 to May 2014. Total 155 patients were included in this study after taking informed consent. An approval was taken from institutional review committee. Patients with hepatic encephalopathy and ascites age from 18 to 65 years were included in this study. Patients with fulminant hepatic failure, non-cirrhotic portal hypertension and uremic, anoxic, cerebral and metabolic encephalopathy were excluded from this study.

Detailed history of all patients were taken and physical examination was done. Patients were labeled as having SBP when ascetic fluid neutrophil count > 250/ml. Hepatic encephalopathy was graded according to west heaven criteria as shown in table 1.

Demographic data of all the patients and clinical and laboratory findings were recorded in pre-designed proforma. All the data was entered in SPSS version 16. Mean and standard was calculated for numerical data. Frequencies were calculated for categorical data and chi-square test was applied to see the association of SBP with grades of HE. P. value ≤ 0.05 was considered as significant.

RESULTS

Total 155 patients with hepatic encephalopathy were included in this cross sectional study. Mean age of patients was 38.9±11.7. Among the 155 patients of hepatic encephalopathy, 119(77%) were males and 36(23%) were females as shown in Figure 1. Among the 155 patients, HBsAg Positive patients were 60(38.70%), Anti-HCV Positive were 87(56.13%) and Both HBsAg & Anti-HCV Positive patients were 8(5.16%) (Table 2). Grading of the patients of hepatic encephalopathy was done according to west heaven criteria (Table 1). Out of 155 patients 48(30.97%) patients had Grade I hepatic encephalopathy and SBP was present in 33 (68.8%) patients, 63 (40.64%) patients had Grade II hepatic encephalopathy and SPB was found in 32(50.8%) patients, 23(14.84%) had Grade III hepatic encephalopathy and SBP was present in 10(43.5%) patients and 21(13.55%) had Grade IV hepatic encephalopathy and SBP was found in 6(28.6%) patients. There is significant association between SBP and grade hepatic encephalopathy (P. value 0.01) (Table 3)

Table 1: West Haven criteria for grading hepatic encephalopathy

Grade 1	Trivial lack of awareness Euphoria or anxiety Shortened attention span Impaired performance of addition
Grade 2	Lethargy or apathy Minimal disorientation for time or place Subtle personality change Inappropriate behavior Impaired performance of subtraction
Grade 3	Somnolence to semi stupor, but responsive to verbal stimuli Confusion Gross disorientation
Grade 4	Coma (unresponsive to verbal or noxious stimuli)

Table 2: Characteristics of patients


Viral cause of cirrhosis HBsAg Positive	60(38.70%)
Anti-HCV Positive	87(56.13%)
Both HBsAg & Anti-HCV Positive	8(5.16%)

Table 3

Grades	Spontaneous Bacterial Peritonitis		Total
	Present	Absent	
Grade I	33(68.8%)	15(31.3%)	48(30.97%)
Grade II	32(50.8%)	31(49.2%)	63(40.64%)
Grade III	10(43.5%)	13(56.5%)	23(14.84%)
Grade IV	6(28.6%)	15 (71.4%)	21(13.55%)
Total	81(52.3%)	74 (47.7%)	155

P value: 0.01

Fig. 1



DISCUSSION

HE is one of the commonest presentation in advanced liver diseases. It precipitates in cirrhosis of liver patients due to SPB, massive ascites with albumin level <1.5g/dl etc. The exact pathogenesis of HE is not well defined, multiple hypothesis are there. In majority of patients the precipitating factor control or reversal is the main corner stone of the HE management¹².

In present study SBP was presented 52.3% patients. This is very high presence of SBP in patients of liver cirrhosis with hepatic encephalopathy. This high percentage of SBP in these patients is due to not taking primary or secondary prophylaxis of antibiotics. SBP needs early diagnosis to prevent mortality in HE patients¹⁰. Ram N et al found 48.46% patients with SBP which is almost close to our study¹⁰. Mumtaz et al is in contrast with this study, who found SBP in 20.5% patients with hepatic encephalopathy¹³. SBP was detected the most common 67% precipitating factor by Devrajani et al¹⁴, this percentage is also in contrast with this study.

Primary prophylaxis of antibiotics is recommended for high risk cirrhotic patients with ascitic fluid albumin <1.5gm/dl⁹. Antibiotic prophylaxis in cirrhotic patients with ascites is mandatory. Variceal bleed is the most common event in advanced cirrhosis. It leads to hepatic encephalopathy and SBP. However antibiotic therapy decrease infection (SBP) and increase survival. In our study, SPB more observed in grade I and II of hepatic encephalopathy. This is due to greater risk factor of GIT bleed in these. Hepatic encephalopathy and SBP both are important complications in cirrhotic patients and potentiate each other in the presence of any participating factors.

CONCLUSION

HE and SBP are the common complications in liver cirrhosis. Most of the HE patients having SBP as well as precipitating factor. The treatment of SBP is the important step of management of HE in liver cirrhosis.

REFERENCES

1. Frederick RT. Current Concepts in the Pathophysiology and Management of Hepatic Encephalopathy. *Gastroenterol Hepatol* (N Y). 2011 Apr;7(4):222–33.
2. Abrams GA, Fallon MB. Cirrhosis of liver and its complications. In: Andreoli TE, Bennett JC, Carpenter CJ, Plum F. editors. *Cecil essentials of medicine*. 5th ed. Philadelphia:WB Saunders; 2001.p.340-5.
3. Assy N, Rosser BG, Grahame GR, Minuk GK. Risk of sedation for upper GI endoscopy exacerbating subclinical hepatic encephalopathy in patients with cirrhosis. *Gastrointest Endosc* 1999; 49:690-4.
4. Ayub A, Ishfaq M. Spontaneous Bacterial Peritonitis; Frequency of asymptomatic in patients of liver cirrhosis. *Professional Med J* 2014;21(4): 606-612.
5. Amin K, Alam MM, Abbas G, Mehmood K, Irshadul-Haq. Spontaneous bacterial peritonitis; prevalence in cirrhotic patients. *Prof Med J* 2000; 7:346-51.
6. Desai AP, Reau N, Reddy KG, Te HS, Mohanty S, Satoskar R, et al. Persistent spontaneous bacterial peritonitis: a common complication in patients with spontaneous bacterial peritonitis and a high score in the model for end-stage liver disease. *Therap Adv Gastroenterol*. 2012 Sep;5(5):275–83.
7. Taneja SK, Dhiman RK. Prevention and Management of Bacterial Infections in Cirrhosis. *International Journal of Hepatology*. 2011 Aug 28;2011:e784540.
8. Pathak R, Harsh S, Adhikari A, Khadga PK. Clinical Spectrum of Spontaneous Bacterial Peritonitis in Tertiary Care Centre. *www jiom com np*. 2013;35:41.
9. Alaniz C, Regal RE. Spontaneous Bacterial Peritonitis. *P T*. 2009 Apr;34(4):204–10.
10. Ram N, Qadri HR, Shah A, Shaikh AS. Frequency of spontaneous bacterial peritonitis in patients with hepatic encephalopathy. *MC*.2012;19(2):36-38.
11. G Garcia-Tsao. Bacterial infections in cirrhosis. *Can J Gastroenterol* 2004;18(6):405-406.
12. Maqsood S, Saleem A, Iqbal A, Butt JA. Precipitating factors of hepatic encephalopathy: experience at Pakistan Institute of Medical Sciences Islamabad. *J Ayub Med Coll Abbottabad* [Internet]. 2006 [cited 2014 Apr 22];18(4). Available from: <http://ayubmed.edu.pk/JAMC/PAST/18-4/Saad%20Masood>.
13. Mumtaz K, Ahmed US, Abid S, Baig N, Hamid S, Jafri W. Precipitating factors and the outcome of hepatic encephalopathy in liver cirrhosis. *J Coll Physicians Surg Pak* 2010; 20: 514-518
14. Devrajani BR, Shah SZA, Devrajani T, Kumar D. Precipitating factors of hepatic encephalopathy at a tertiary care hospital Jamshoro, Hyderabad. *J Pak Med Assoc*. 2009 Oct;59(10):683–6.