

# Complications in Patients of Acute/Chronic Renal Failure during Haemodialysis

NAZIR AHMAD<sup>1</sup>, KASHIF SARDAR<sup>2</sup>, MOHAMMAD HUMAYUN<sup>3</sup>

## ABSTRACT

**Aim:** To determine the frequency of different complications during haemodialysis.

**Study design:** Cross sectional study.

**Place and duration:** This study was conducted at District Teaching Hospital, Ghazi Khan Medical College, DG Khan from January 2014 to December 2014. A total of 227 patients were included.

**Methods:** The study was conducted in Dialysis unit of District Teaching Hospital, Ghazi Khan Medical College, DG Khan. 227 patients were selected who were undergoing haemodialysis due to acute/chronic renal failure. All the patients were evaluated for possible complications of haemodialysis namely psychological/GIT/cardiovascular and Neurological complications.

**Results:** The most common complication was depressive illness and was observed in 140 (61.6%) patients, other vomiting, fever, itching and hypotension. The causative relations of the complications with other parameters were assessed.

**Conclusion:** A very high frequency of the complications associated with maintenance haemodialysis.

**Key words:** Acute renal failure, chronic renal failure, haemodialysis

---

## INTRODUCTION

Chronic renal failure may be defined as a glomerular filtration rate (GFR) below 30ml per minute<sup>1</sup>. Symptoms and complications of uremia often occur when GFR is <15 ml per minute. Chronic renal failure affects every aspect of life of the patients who suffer from it. Over 165000 patients in United States were treated for end stage renal disease during 1990<sup>2</sup>. Patients with end stage renal disease requires replacement treatment. This can be provided by maintenance haemodialysis, peritoneal dialysis or kidney transplantation. In Europe, over 250 per million populations are given replacement treatment for end stage renal failure. Only about 30% of these patients undergo renal transplantation, which is considered as the gold standard of renal replacement therapy<sup>3</sup>.

The dialysis is only a part of end stage renal management because it cannot replace all the improvement functions of kidney. During haemodialysis the patient may develop complications, e.g., many factors like size of extracorporeal circulation, degree of ultra-filtration, changes in serum osmolality, autonomic neuropathy and alteration in body temperature due to dialysis temperature<sup>4</sup>. Vomiting is another common complication during dialysis.

The clinical features include shivering with anaphylactic reaction of varying intensity from mild respiratory distress to anaphylactic shock. It is because of substance released by extra-corporeal circuit and some patients were thought to be intolerant of dialysis membrane made up of cuprophane, which may be due to brisk complement activation<sup>5</sup>.

Disequilibrium syndrome is due to rapid flux in osmolality, with haemodialysis. The clinical features include confusion, clouding of consciousness and fits. Mechanical and estrogenic complications are less common with improved equipment. These complications are air embolism, haematemeses, blood and contaminated dialysate. Complications related to vascular access problems are infection at the site of AV fistulae<sup>6</sup> or infection due to subclavian or femoral vein catheterization, stenosis of fistula, haemorrhage, thrombosis, inflammatory complications, aneurysm and undesirable consequences of fistula.

## MATERIALS AND METHODS

This cross sectional study was conducted at Dialysis Centre District Teaching Hospital, Ghazi Khan Medical College, DG Khan from July 2012 to January 2013. A total of 227 patients were included in the study. Patients suffering from acute/chronic renal illness and coming for dialysis to Dialysis Unit of District Teaching Hospital, Ghazi Khan Medical College, DG Khan were included. General population and refusal to participate were excluded from the study.

---

<sup>1</sup>Assistant Professor of Medicine, District Teaching Hospital, Ghazi Khan Medical College, DG Khan,

<sup>2</sup>Assistant Professor of Medicine, Multan Medical & Dental College

<sup>3</sup>Associate Professor/Head of Medical Entomology & Parasitology Department, Institute of Public Health, Lahore

Correspondence: Dr. Nazir Ahmad, e-mail: drmalik.nazir.ahmed@gmail.com

**RESULTS**

Out of 227 patients, 123(54.2%) were male and 104 (45.8%) were female. Mean age of the patients was 33.3±10.5 ranging from 13-65 years. Psychological complications were observed in 180 patients. The most common complication was depressive illness was observed in 140(61.6%) [80 males and 60 females] patients. Impotence was found in 40 (17.8%) patients. Shivering was found in 59(57.3%) males and 44(42.7%) female patients whereas fever was seen in 63(62.4%) males and 38(37.6%) females (Tables 1-3).

Table 1: GIT related complications

Complication	Male	Female	Total
Vomiting	70(30.8)	39(17.2)	109(48%)
Pain abdomen	04(1.76%)	06 (2.6%)	10(4.5%)
Nausea	03(1.32%)	02 (.88%)	05(2.2%)
Haematemesis	03(1.32%)	01 (.44%)	04(1.5%)

Table 2: Cardiovascular complications

Complication	Male	Female	Total
Cardiac arrhythmias	110(49%)	60 (26%)	170(75%)
Hypokalemia	36(15.8%)	20(8.8%)	56(25%)
Hypertension	10 (4.4%)	06 (2.6%)	16(7%)
Air embolism	07 (3.1%)	4(1.9%)	11(5%)

Table 3: Neurological complications

Complication	Male	Female	Total
Headache	5 (2.2%)	9 (3.9%)	14(6.2%)
Fits	3(1.32%)	5 (2.2%)	08(3.5%)
Disequilibrium	4 (1.5%)	3(1.32%)	06(2.6%)
Dementia	3(1.32%)	2 (.88%)	05(2.2%)

**DISCUSSION**

The major renal replacement therapy is haemodialysis worldwide in the management of end stage renal diseases<sup>8,9</sup>. It is either sole replacement therapy or prior to renal transplantation.

The commonly observed GIT related complications during haemodialysis are vomiting, pain in abdomen and haematemesis, 48% of the patients had vomiting during or after dialysis. These results are comparable to the results of a study<sup>8</sup>. Heparinization during haemodialysis can precipitate upper gastrointestinal bleeding responsible for haematemesis, though it is not a very common problem. In present study, 4 patients during study period experienced the upper GI bleeding. International data are also documenting the low incidence of haematemesis as the complication of maintenance haemodialysis.<sup>9</sup>

In present study the complications of the haemodialysis occurred in 80% of the patients, either

a major one or a minor. The psychological complications were at the top with reference to the frequency. These were observed in 79% of the patients. The most common psychological complication was depressive illness, occurring in 61% of the dialyzed patients, which is in line to that on other studies<sup>10,11</sup>.

The cardiovascular complications, like hypertension, arrhythmia have also been observed, hypokalemia being the cause of arrhythmia. It requires termination of dialysis until and unless the management of hypokalemia. Hypokalemia, hypertension and air embolism were observed in 25%, 7% and 5% respectively. It has been suggested that hypotension and associated symptoms may be related to the biochemical changes caused by cellulosic membranes or due to hypovolemia or excessive ultrafiltration of water at the time of dialysis<sup>10</sup>. The ultrafiltration depends upon the condition of the patient, if hypotension and hypovolemia is evident, then no ultrafiltration is advised.

Cardiac arrhythmias are frequently associated with haemodialysis and have been reported to be seen in some series among 76% of the patients<sup>11</sup>. These appeared to occur mostly after the first three hours of dialysis. Various risk factors predisposing dialysis patients to arrhythmias have been identified including rapid electrolyte and fluid shifts, acid base alterations, alteration in parathyroid and calcium metabolism, reduced oxygen saturation levels and changes in erythrocyte's potassium levels<sup>12</sup>.

Clinicians have long observed that male patients with haemodialysis have the problem of impotence. A study was conducted on 32 male haemodialyzed patients. The views of all these patients were separately interviewed. The reported frequency of impotence was 38%<sup>12</sup>. In another study the prevalence of impotence in men on haemodialysis in excess of 70% was reported. In this study 17.8% of male patients reported the impotence.

Headache, fits, disequilibrium and dementia were the neurological complications observed in 6%, 5%, 4% and 3% respectively. These complications in this study also reflect low frequency in contrast to other international studies<sup>13,14</sup>. This contrast is due to the short duration of maintenance dialysis in this study as compared to international data in which patients were on maintenance haemodialysis for the period of more than 5 years.

The shivering, fever and itching were the other common recorded complications of haemodialysis. These were observed in 45%, 42% and 22% respectively. These results are comparable with other studies<sup>10,15</sup>.

## CONCLUSION

The results of this study revealed a very high frequency of the complications associated with maintenance haemodialysis.

## REFERENCES

1. Walls J. Chronic renal failure: causes and conservative management. *Med Intern* 1995; 9(30): 144-8.
2. Warnock DG. Chronic renal failure. In: Bunnet editor. *Cecil textbook of medicine*. 20<sup>th</sup> ed. Philadelphia: WB Saunders, 1996; 1: 556-63.
3. Kishimoto TK, Viswanathan K, Ganguly T. Contaminated heparin associated with adverse clinical events and activation of the contact system. *N Engl J Med* 2008; 358(23): 2457-67.
4. Weinreich T, De los Ríos T, Gaulty A, Passlick-Deetjen J. Effects of an increase in time vs. frequency on cardiovascular parameters in chronic hemodialysis patients. *Clin Nephrol* 2006; 6(6): 433-9.
5. Eknoyan G, Beck GJ, Cheung AK. Effect of dialysis dose and membrane flux in maintenance hemodialysis. *N Engl J Med* 2002; 347 (25): 2010-9.
6. Cheung AK, Levin NW, Greene. Effect of high flux hemodialysis on clinical outcomes. *J Am Soc Nephrol* 2003; 14(12): 3251-63.
7. Locatelli F, Martin-Malo A, Hannedouche T. Effect of membrane permeability on survival of hemodialysis patients. *J Am Soc Nephrol* 2009; 20 (3): 645-54.
8. van rassel MS, Koppius PW, Schut NH. Headache during hemodialysis. *Clin Nephrol* 2008; 69: 219.
9. Kapa S, Qian Q. 84 year old woman with hemodialysis-associated shortness of breath. *Mayo Clin Proc* 2009; 84: 187.
10. Kitano Y, Kasuga H, Watanabe MI. Severe coronary stenosis is an important factor for induction and lengthy persistence of ventricular arrhythmia during and after hemodialysis. *Am J Kid Dis* 2004; 44: 328.
11. Weiner DE, Nicholls AJ, Sarnak MJ. Cardiovascular disease. In: *Handbook of Dialysis*, Daugirdas JT, Blake PG, Ing TS (Eds), Lippincott, Williams & Wilkins, Philadelphia 2007. p.626.
12. Grebenyuk LA, Marcus RJ, Nahum E. Pulmonary embolism following successful thrombectomy of an arteriovenous dialysis fistula. *J Vasc Assess* 2009; 10: 59.
13. 13. Miljkovic M, Zidverc TJ, Stemic N. Hemodialysis headache. *Clin Nephrol* 2009; 71: 158.
14. Buemi N, Coppolino G, Boligmamp D. Arrhythmias and hemodialysis. *Ren Fail* 2009; 31:75.
15. Kamik JA, Young BS, Lew N. Cardiac arrest and sudden death in dialysis units. *Kid Int* 2001; 60: 350.