

Evaluation of Serum Cholesterol and Triglycerides in Patients with Diabetes and Hypertension

MUHAMMAD ASLAM SOOMRO¹, AZEEM IQBAL², MUHAMMAD MOHSIN SARDAR³

ABSTRACT

Aim: To determine the percentage of hypertensive and diabetic patients having increased level of cholesterol and triglycerides.

Methods: This descriptive multicentrestudy was carried out on cases of diabetes mellitus, hypertension and glycemic control from 1st June 2016 at 31st December 2016. A total of 300 cases were included in the study.

Results: Out of 300 patients, 114(38%) showed isolated diabetes mellitus. Isolated hypertension was found in 21(7%), 6(2%) patients showed combined hypertension and diabetes mellitus. 18(6%) patients exhibited combined hypertension and stroke. 20(6.7%) patients were having combined diabetes mellitus and hypertension and 7(2.3%) patients exhibited diabetes mellitus.

Conclusion: Dyslipidemia is significantly higher in diabetic and hypertensive population compared to controls.

Keywords: Diabetes mellitus, Hypertension

INTRODUCTION

Cholesterol is a fatty material, which over the years thickens and narrows the arterioles. It is manufactured in the liver and is an essential element of blood. It is present abundantly in egg yolk, butter, cream, animal fat and milk. Hypertensive retinopathy is not only the result of elevated blood pressure, although it is established that increased blood pressure caused failure of the auto-regulation of retinal circulation.¹

The major risk factors contributing to the excess of cardiovascular disease caused by diabetes includes hyperglycemia, insulin resistance, dyslipidemia, hypertension and smoking. Dyslipidemia in hypertensive patients is itself known to be a predisposing risk factor, an aggravating or complicating factor². Hypertension and hyperlipidemia not only accelerate atherogenesis but also cause degenerative changes in the walls of large- and medium-sized arteries³, which accelerate cerebrovascular hemorrhage⁴, ischemic heart disease⁵, stroke, and cardiac arrest^{6,7,8}.

Increased serum cholesterol levels have been observed in individuals who suffer from non-haemorrhagic stroke, but not in those who suffer intra cranial or subarachnoid haemorrhage. Serum lipids are thought to interact with the pathogenesis of non-haemorrhagic stroke through atherosclerosis mechanism.

¹Associate Prof Medicine, GMMMC, Sukkur/SMBBMU Larkana,
^{2,3}HO Internal Medicine, ShaikhZayed Hospital, Lahore
Correspondence Dr. Muhammad AslamSoomro
Email: dr_soomro_aslam@outlook.com

PATIENTS AND METHODS

This descriptive multicentrestudy was carried out at Civil Hospital, GMMMC Sukkur and ShaikhZayed Hospital Lahore cases of diabetes mellitus, hypertension and glycemic control from 1st June 2016 at 31st December 2016. A total of 300 cases were included in the study.

RESULTS

Out of 300 patients, 153(51%) were male and 147(49%) were female. Hypertriglyceridemia was present in 18(46%) male and 21(54%) in female patients whereas hypercholesterolemia was seen in 9(60%) male and 6(40%) in female patients.

Table1: Triglycerides and cholesterol level in patients with hypertension (n=42)

Age (yrs)	Male	Female
< 40	9	03
41-50	-	-
51-60	9	6
61-70	3	9
>70	3	-

Table2: Triglycerides and cholesterol level in patients with diabetes (n=114)

Age (yrs)	Male	Female
< 40	39	15
41-50	9	09
51-60	33	27
61-70	9	09
>70	-	-

Dyslipidemia in hypertension patients, hypertriglyceridemia was present in 12(67%) male and (33%) in female patients and hypercholesterolemia was seen in 6(67%) male and 3(43%) in female patients. Dyslipidemia in diabetic patients, hypertriglyceridemia was present in 15(38%) male and 24(68%) in female patients, hypercholesterolemia was seen in 12(50%) male and 12(50%) in female patients.

DISCUSSION

In the present study 114(38%) patients of the total 300 patients with mean age of 54 years were suffering from isolated diabetes mellitus. Of these 114 patients, 45% were male while 55% were female. In this study 21% of the patients were having increased level of serum cholesterol and 34.2% patients having increased level of triglyceride (>150mg/dl). When we compare this study with a study, it was found that 24% of the population was smoker and 64% of the patients were having sedentary life style⁸.

Another study showed that the effect of flavastatin HMG CoA reductase inhibitor was assessed in 50 patients with dyslipidemia in diabetic belonged to the city of Lahore.⁹ In this study the total cholesterol and triglyceride were high. When compare this study with the present one, total cholesterol and triglycerides are quite low in present study. The reason is financial status, difference between people belonging to Lahore and our results are comparable.

Hyperglyceridemia is most common lipid abnormality in NIDDM, particularly in diabetics with poor glycemic control.¹⁰ The above observation was also seen in present study. Triglyceride levels were above the desired level in 34.2% patients out of 114 patients, with mean triglyceride level below 150mg/dl. Hyperglyceridemia is a potent risk factor for macrovascular disease due to reduced synthesis of insulin dependent lipoprotein lipase in liver, resulting in impaired clearance of LDL, chylomicron and VLDL remnants¹¹. High triglyceride levels are associated with hypercoagulopathy and decreased fibrinolysis both contributing to coronary heart disease¹². In addition to metabolic relation to LDL cholesterol modifying its particle into more dense, small and hence more atherogenic form¹³.

In the present study 42(14%) patients with mean age of 63 years were suffering from isolated hypertension. Of these 57% were male while 43% had increased level of triglyceride and 22% were having increased serum cholesterol. In these patients 64% were smoker. The similar study showed that 32% of the patients were having

hyperglyceridemia and 10% of patients were having both raised triglyceride and cholesterol¹⁴.

In our study more patients had increased level of cholesterol and triglyceride. The exact cause of increased cholesterol and triglyceride in present study is not known. A study carried out reveals that 48% of patients were having hypercholesterolemia¹⁵. Another study showed that 50% of the patients had dyslipidemia and 39% had hypertriglyceridemia¹⁶.

A study showed the frequent occurrence of high serum lipids in hypertensives which has been termed as familial dyslipidemic hypertensive.¹⁷ Hypertriglyceridemia was the most prevalent of all the lipid abnormalities^{18,19}.

CONCLUSION

Although the sex ratios may vary, dyslipidemia is found in significant number of diabetics and hypertensives as compared to controls.

REFERENCES

1. Liebreich R. Bright's disease at diagnosis-shear ophthalmoscopy Albercht Von Graefes. Arch Ophthalmol 2007; 85: 5:265-8.
2. Thakur AK, Achari V. A study of lipid levels in uncomplicated hypertension. Indian Heart J 2000; 52: 173-7.
3. Naomi DL, Gordon HW. Hypertensive Vascular Disease. In: Fisher ND, Williams GH, editors. Harrison's Principles of Internal Medicine. 16th Ed. New York: McGraw Hill; 2005. p. 1463-70.
5. Marshal SM, Flvbjerg A. Prevention and early detection of vascular complications of diabetes. BMJ 2006; 333(7566): 475-80.
6. Buse JB, Bigger JT, Bvington RP. Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial: design and methods. Am J Cardiol 2007; 99(12A):21i-33i.
7. Hemmingsen B, Lund SS, Gluud C, Vaag A, Almdal TP. Targeting intensive glycaemic control versus targeting conventional glycaemic control for type 2 diabetes mellitus. Cochrane Database Syst Rev. 2013; 11.:CD008143.
8. Duncan BB, Wong TY, Tyroler HA, Davis CE, Fuchs FD. Hypertensive retinopathy and incident coronary heart disease in high risk men. Br J Ophthalmol 2002; 86: 1002-6.
9. Ye J, Kiage JN, Arnett DK, Bartolucci AA, Kabagambe EK. Short-term effect of fenofibrate on C-reactive protein. Diabetol Metab Syndr 2011; 3: 24.
10. Kurozumi A, Okada Y, Mori H, Arai T, Tanaka Y. Efficacy of α -glucosidase inhibitors combined with dipeptidyl-peptidase-4 inhibitor. J Diabetes Invest 2013; 4(4): 393-8.
11. Ferreira M, Ant3nio N, Gonalves F. Hemoglobin: simply a laboratory value or a powerful predictor of

- risk in patients with acute coronary syndrome?. *Rev Port Cardiol* 2012; 31(2): 121-31.
12. Hui ML, Kumar A, Adams GG. Protocol-directed insulin infusion sliding scales improve perioperative hyperglycaemia in critical care. *Perioper Med (Lond)* 2012; 1: 7.
 13. Kuwashiro T, Sugimori H, Ago T, Kamouchi M, Kitazono T. Risk factors predisposing to stroke recurrence within one year of non-cardioembolic stroke onset: the Fukuoka Stroke Registry. *Cerebrovasc Dis* 2012; 33(2): 141-9.
 14. Ikeda N, Iijima R, Hara H, Moroi M, Nakamura M, Sugi K. Glycated hemoglobin is associated with the complexity of coronary artery disease, even in non-diabetic adults. *J AtherosclerThromb* 2012; 19(12):1066-72.
 15. Stenestrand U, Wijkman M, Fredrikson M, Nystrom FH. Association between admission supine systolic blood pressure and 1-year mortality in patients admitted to the intensive care unit for acute chest pain. *JAMA* 2010; 303(12): 1167-72.
 16. Sandercock P, Mielke O, Liu M, Counsell C. Anticoagulants for preventing recurrence following presumed non-cardioembolic ischaemic stroke or transient ischaemic attack. *Cochrane Database Syst Rev* 2003; (1): CD000248.
 17. Wu S, Shi Y, Wang C. Glycated hemoglobin independently predicts stroke recurrence within one year after acute first-ever non-cardioembolic strokes onset in A Chinese cohort study. *PLoS One* 2013; 8(11): e80690.
 18. Thakur AK, Achan V. A study of lipid levels in uncomplicated hypertension. *Heart J* 2000; 52: 173-7.
 19. Rejmohan I, Deepa R, Mosan A, Mohan V. Association between isolated hypercholesterolemia. *Heart J* 2000; 52: 400-6.