

Evaluation of Hypovitaminosis D in different age and gender groups in Khyber Pakhtunkhwa Pakistan

SAJID ALI¹, MUHAMMAD ZUBAIR², AHMAD SHAMIM KHAN³, SIDRA ZAMAN⁴, ALISHBA ALI⁴, SAWERA IFTIKHAR⁴¹Associate Professor, Department of Biotechnology, Abdul Wali Khan University, Mardan, Pakistan.²Registrar, Department of Nephrology, Dubai Health Authority, Dubai.³Registrar, Department of Nephrology, Pakistan Institute of Medical Sciences, Islamabad, Pakistan.⁴Bachelor Students, Department of Biotechnology, Abdul Wali Khan University, Mardan, Pakistan.Corresponding author*: Dr. Sajid Ali, Email: sajid@awakum.edu.pk

ABSTRACT

Background: Hypovitaminosis D is associated with many chronic disorders such as cancer, osteoporosis, cardiovascular, rickets, and musculoskeletal disorders. Therefore, this study was designed to determine the level of vitamin D in our region.

Methods: A total of one hundred and ninety-five adult participants were enrolled. Patients were referred by medical physicians and consultants after identifying the sign and symptoms of Hypovitaminosis D. Both male and female gender were recruited and patients with severe chronic disorders were excluded from the study. Vitamin D levels were categorized into four groups including deficiency of vitamin D (<10 IU), insufficient vitamin D level (10-30 IU), Normal vitamin D level (30-100 IU), and Hypervitaminosis level of vitamin D (>100 IU).

Results: The minimum and maximum age of the patient was 17 and 57 years age with the mean age of all patients being 33.833 ± 17.3 . Male and female patients were 79.5% (n=155) and 20.5% (n=40). The highest number of patients were identified in the age group 36-45 years. Vitamin D level was found in 7.7% deficient while insufficiency of vitamin D was found in 48.7%. Insufficiency of vitamin D was found more in female (67.5%) patients than in males (43.9%). The highest prevalence of vitamin D insufficiency was obtained in the age group 16-25 years. No patient was found with hypervitaminosis D.

Conclusion: There is a high deficiency of vitamin D in patients particularly in females and younger age populations. Awareness about vitamin D importance, psychosocial intervention, and vitamin D supplementation should be implemented to prevent any severe consequences of Hypovitaminosis D.

Keywords: Vitamin D, Hypovitaminosis D, Age, Gender

INTRODUCTION

Vitamin D is also called sunshine vitamin or also referred to as calciferol. Vitamin D insufficiency affects almost 50% of the population worldwide¹. Hypovitaminosis D is a pandemic in which several factors contribute including environmental, and lifestyle². Hypovitaminosis D is an important public health problem³. Vitamin D play essential role in cell proliferation, differentiation, and apoptosis⁴. Hypovitaminosis D may leads to rickets in children due to bone softening and de-mineralization^{5,6}. While in adults particularly in women cause osteomalacia and osteoporosis, resulting of bone fractures^{7,8}.

Vitamin D receptors are distributed throughout the body and they have a role in greater than 200 human genes⁹, showing diverse functions in maintaining human health¹⁰. Vitamin D has the main role in immunity, reproductive, muscular, skeletal, and integumentary systems in all ages of people¹¹. It is reported that vitamin D level is significantly associated with chronic diseases¹².

Several factors are involved in Hypovitaminosis D such as old age, race, use of medications having antagonist effects on the metabolism of vitamin D, high body mass index, inadequate use of vitamin D supplementation, lack of exercise, education, and awareness about vitamin D importance¹³. Presently, Hypovitaminosis D has been reported pandemic throughout the world^{1,14} and reported 30-93% deficiency in different studies^{15,16}.

For the last two decades, it is reported that Hypovitaminosis D is also high in tropical countries including China¹⁷, India¹⁸, Iran¹⁹, Saudi Arabia²⁰, and Turkey²¹ where sunlight plays a significant role. Pakistan is considered one of the sun-drenched countries in the globe and its supposed to have enough sunlight to maintain an adequate level of vitamin D but Hypovitaminosis D is also prevalent in Pakistan²². Hypovitaminosis D is also reported in pregnant²³, lactating women, and infants in Pakistan^{24,25}. Few studies reported from our region therefore, this study was designed to determine the level of vitamin D in our population.

MATERIALS AND METHODS

This descriptive cross-sectional study was conducted in a duration of three months from January to March 2022. The study was carried out in the department of Biotechnology at Abdul Wali Khan University, Mardan, along with the collaboration of Real-Time PCR

Laboratory, Dabgeri Garden, Peshawar, Pakistan. One hundred and ninety-five patients were recruited for the present study. All those patients who had been referred by medical physicians and consultants based on the sign and symptoms of Hypovitaminosis D in patients were included irrespective of age, gender, and ethnicity of patients. Whereas those patients had other comorbidities including celiac disease, unexplained musculoskeletal pain, cystic fibrosis, bony deformities or tenderness, Crohn's disease, metabolic bone disease, patients had a history of any surgery on the intestine or stomach, malignancy, patients with a history of pituitary gland problem, pregnant and lactating females, kidney diseases, asthmatic patients and other known chronic disease patients were excluded from the study. Moreover, those patients who were not willing to take participation in the research study were excluded from the present study.

Ethical approval was taken from the Real-Time PCR Laboratory, Dabgeri Garden, Peshawar, and the Department of Biotechnology at Abdul Wali Khan University, Mardan. Informed consent was obtained from the patients before recruitment in the study.

Vitamin D levels were evaluated through the detection of 25-OH-D levels. Therefore, 5 ml peripheral blood was collected in a serum separator tube (SST provided by BD vacutanier®) from all the participants at the time of the laboratory visit after the medical physician and consultant checkup. The collected sample was centrifuged and the serum 25-OH-D level was determined through the Enzyme-linked immunoassay technique (ELISA). The participants were categorized into four groups according to 25-OH-D levels in their sera: 1) Deficiency: <10 IU; 2) Insufficient: 11-30 IU; 3) Normal: 31-100 IU; and 4) Hypervitaminosis D/Toxicity >100 IU.

Descriptive statistics were used for demographic, and laboratory test data, and proportions were observed in collected data. The data were analyzed through Statistical Package for Social Science (SPSS) version 22. Means and standard deviation were determined.

RESULTS

A total of one hundred and ninety-five participants (n=195) were included. The mean age of patients was 33.8 ± 17.3 with a minimum

of 17 years of age and a maximum of 57 years of age. Among the total, 79.5% (n=155) were male patients with a mean age of 35.3 ± 10.1 . Whereas female patients were 20.5% (n=40) with a mean age of 28.7 ± 9.91 . All patients were categorized into five groups. The highest number of participants were determined in the age group 36-45 years (29.2%), followed by the age group 26-35 years (28.7%), 16-25 years age group (27.7%), 46-55 years age group (11.3%), and 55-65 years age group (3.1%) (Table 1).

Table 1: Distribution of study participants in different Age and Gender groups (n=195)

Age (Years)	Male % (n)	Female % (n)	Total % (n)
16-25	20.6 (32)	55.0 (22)	27.7 (54)
26-35	30.3 (47)	22.5 (09)	28.7 (56)
36-45	32.9 (51)	15.0 (06)	29.2 (57)
46-55	12.9 (20)	5.0 (02)	11.3 (22)
55-65	3.2 (05)	2.5 (01)	3.1 (06)
Grand Total	79.5 (155)	20.5 (40)	100 (195)

All patients' blood was analyzed for vitamin D levels. Vitamin D levels were categorized into four groups including deficiency of vitamin D (<10 IU), insufficient vitamin D level (10-30 IU), Normal vitamin D level (30-100 IU), and Hypervitaminosis level of vitamin D (>100 IU). Vitamin D level was deficient (7.7%), insufficient (48.7%), and normal (43.6%) in n=15, n=95, and n=85 respectively. No participant was found with hypervitaminosis D. Insufficient level of vitamin D was determined in female (67.5%) patients as compared to males (43.9%) (Table 2).

Table 2: Distribution of Vitamin D levels in males and females (n=195)

Vitamin D Level	Gender Groups		
	Male % (n)	Female % (n)	Total % (n)
Deficiency <10 IU	7.7 (12)	7.5 (03)	7.7 (15)
Insufficient 10-30 IU	43.9 (68)	67.5 (27)	48.7 (95)
Normal 30-100 IU	48.4 (75)	25.0 (10)	43.6 (85)
Hypervitaminosis D/Toxicity >100 IU	0 (0)	0 (0)	0 (0)
Grand Total	79.5 (155)	20.5 (40)	100 (195)

Vitamin D level was also determined in different age groups. The highest prevalence of vitamin D insufficiency was obtained in the age group 16-25 years, followed by the age group 36-45 years, and then the 26-35 years age group. Hypovitaminosis D was found more in the age group 36-45 years (Table 3).

Table 3: Distribution of Vitamin D level in different age groups (n=195)

Vitamin D Level	Age Groups (Years)					
	16-25 % (n)	26-35 % (n)	36-45 % (n)	46-55 % (n)	55-65 % (n)	Total % (n)
Deficiency <10 IU	02	02	11	0	0	7.7 (15)
Insufficient 10-30 IU	45	16	31	02	01	48.7 (95)
Normal 30-100 IU	04	39	15	22	05	43.6 (85)

DISCUSSION

Hypovitaminosis D is a reported pandemic both in developed and underdeveloped countries. Sunlight is a major source of vitamin D and only a few natural foods are enriched with vitamin D. Hypovitaminosis D is more frequently found in Middle East and South Asian countries²⁶. Hypovitaminosis D affects more than 50% population throughout the world and is considered an important public health issue, particularly in developing countries including Pakistan²⁷.

In our study, 56.4% of individuals suffered from low vitamin D levels (Deficiency and insufficient). Riaz et al. published their data of high sample size (n=4830), in which findings were consistent with this study finding that 53.5% had Hypovitaminosis D and 31.5% were insufficient with vitamin D²⁸. Our reported prevalence is much lower than the study reported in which 90% of healthy volunteers were affected by the low level of vitamin D (insufficiency and deficiency)²². Similar to the present study findings, Chaudhary

et al., also revealed that females are more susceptible to Hypovitaminosis D as compared to males²⁹.

In the present study, all patients were categorized into different strata in which the most common Hypovitaminosis D was found younger age group individuals. Similar to our findings, Chaudhary et al., also found that younger age group individuals than 50 years are frequently vulnerable to Hypovitaminosis D than older age individuals²⁹.

A study from Afghanistan reported a significant relationship between gender differentiation and level of vitamin D³⁰. A report from Lahore also revealed an 81% Hypovitaminosis D in females¹⁵. Present reported proportion is lower than the proportion reported by Zargar et al., from Kashmir, India which shows an 83% deficiency of vitamin D³¹. Hypovitaminosis D is more in female and young age people as compared to male and old age people. This Hypovitaminosis D is a great risk which further leads to severe complications. Similar to our study, Azizi et al., from Afghanistan reported that females are more susceptible to Hypovitaminosis D than male individuals³⁰. Similarly, other studies also found like our findings from regions³² of the world³³. Studies show from Kashmir and other regions³⁴ of the world show greater prevalence in females^{31, 35}, which could be explained by less exposure to sunlight, decreased body area exposure, and fewer outdoor activities than male individuals²². A higher prevalence of Hypovitaminosis D may be due to ignoring sunlight due to religious factors (to cover the whole body), fear of darkening of the skin, and traditionally to expose hands and face in the female population for outdoor visits³⁶. Another study reported that sunscreen and heavy clothes/fabrics usage are significantly associated with Hypovitaminosis D³⁷.

The prevalence of Hypovitaminosis D prevalent in male and female are contradictory throughout the world. On the other side, some studies show that vitamin D concentration are found greater in women as compared to men³⁸. Moreover, a study from Saudi Arabia reported that Hypovitaminosis D was found in 70% of male populations as compared to females³⁹. A study from the United States (US) revealed that Hypovitaminosis D is higher in males individuals than females patients^{40, 41}. Which is explained by a higher body mass index, and less consumption of milk²². Additionally, the male subjects also avoid sunlight exposure due to misconceptions related to the harmful effects of sunlight, high temperatures in summer in some parts of the world, or lack of awareness regarding the role of vitamin D in body growth and metabolism³⁶.

In the present study, younger individuals are more vulnerable to Hypovitaminosis D, which may be explained by less consumption of vitamin D-containing foods including oily fish and fortified cereals. Hypovitaminosis D in Pakistan like the developing country is more prevalent due to insufficient dietary supplemented foodstuffs, resulting in low dietary intake of vitamin D²². Hypovitaminosis D have several long term consequences including the diabetic nephropathy⁴².

Another hypothesis shows that air pollution is another factor that prevents ultraviolet exposure to the skin due to the high level of troposphere ozone. Pakistan is one of the polluted countries, therefore pollution could be another factor⁴³. In a few studies, the high percentage of Hypovitaminosis D was attributed to dietary factors^{18, 21}, air pollution⁴⁴, clothing^{21, 45}, and avoidance of sunlight exposure⁴⁶.

The study was conducted on small scale and was a cross-sectional study including only approximately 20% female population with a convenient sampling technique. Blood samples were collected at a single point time and no follow-ups were collected, due to limited financial status, therefore no information was gathered regarding the long-term effects of Hypovitaminosis D. Daily intake of Vitamin D was also not determined due to several reasons. Seasonal/overcast and rainy days can affect vitamin D levels which were not studied. Lifestyle factors and serum parathormone levels were also not checked due to limited resources. Control group and comparison between urban and rural areas people is required to understand the scenario of Hypovitaminosis D.

CONCLUSION

This study shows a higher proportion of Hypovitaminosis D in female and younger age individuals. This increases the risk of developing chronic conditions associated with immunity, and reproductive and musculoskeletal systems. It is important to educate the public regarding the vitamin D role in our health, to avoid any severe complications. A suitable lifestyle and optimum sunlight exposure can prevent Hypovitaminosis D levels. Hence clinical trials and interventional studies are highly necessary to obtain appropriate levels, doses, and affect of supplements of vitamin D. To combat Hypovitaminosis D, it is highly crucial to intake vitamin D-rich food at a high level, increase sunlight exposure, and start food fortification enriched with level of vitamin D.

REFERENCES

- Holick MF. Vitamin D deficiency. *New England journal of medicine*. 2007;357(3):266-81.
- Rostand SG. Ultraviolet light may contribute to geographic and racial blood pressure differences. *Hypertension*. 1997;30(2):150-6.
- Melamed ML, Michos ED, Post W, Astor B. 25-hydroxyvitamin D levels and the risk of mortality in the general population. *Archives of internal medicine*. 2008;168(15):1629-37.
- Thomas MK, Lloyd-Jones DM, Thadhani RI, Shaw AC, Deraska DJ, Kitch BT, et al. Hypovitaminosis D in medical inpatients. *New England Journal of Medicine*. 1998;338(12):777-83.
- Holick MF, editor. High prevalence of vitamin D inadequacy and implications for health. *Mayo Clinic Proceedings*; 2006: Elsevier.
- Gordon CM, Williams AL, Feldman HA, May J, Sinclair L, Vasquez A, et al. Treatment of hypovitaminosis D in infants and toddlers. *The Journal of Clinical Endocrinology & Metabolism*. 2008;93(7):2716-21.
- Bischoff-Ferrari HA, Willett WC, Wong JB, Giovannucci E, Dietrich T, Dawson-Hughes B. Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *Jama*. 2005;293(18):2257-64.
- Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, Orav JE, Stuck A, Theiler R, et al. Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *Bmj*. 2009;339.
- Bouillon R, Marcucci C, Carmeliet G, Bikle D, White JH, Dawson-Hughes B, et al. Skeletal and extraskeletal actions of vitamin D: current evidence and outstanding questions. *Endocrine reviews*. 2019;40(4):1109-51.
- Ramagopalan SV, Heger A, Berlanga AJ, Mauger NJ, Lincoln MR, Burrell A, et al. A ChIP-seq defined genome-wide map of vitamin D receptor binding: associations with disease and evolution. *Genome research*. 2010;20(10):1352-60.
- Whiting SJ, Calvo MS. Dietary recommendations for vitamin D: a critical need for functional end points to establish an estimated average requirement. *The Journal of nutrition*. 2005;135(2):304-9.
- Heaney RP. Long-latency deficiency disease: insights from calcium and vitamin D. *The American journal of clinical nutrition*. 2003;78(5):912-9.
- Holick MF, Siris ES, Binkley N, Beard MK, Khan A, Katzer JT, et al. Prevalence of vitamin D inadequacy among postmenopausal North American women receiving osteoporosis therapy. *The Journal of Clinical Endocrinology & Metabolism*. 2005;90(6):3215-24.
- Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *The American journal of clinical nutrition*. 2008;87(4):1080S-6S.
- Masud F. Vitamin D levels for optimum bone health. *Singapore medical journal*. 2007;48(3):207.
- Isaia G, Giorgino R, Rini G, Bevilacqua M, Maugeri D, Adami S. Prevalence of hypovitaminosis D in elderly women in Italy: clinical consequences and risk factors. *Osteoporosis International*. 2003;14(7):577-82.
- Du X, Greenfield H, Fraser DR, Ge K, Trube A, Wang Y. Vitamin D deficiency and associated factors in adolescent girls in Beijing. *The American journal of clinical nutrition*. 2001;74(4):494-500.
- Goswami R, Gupta N, Goswami D, Marwaha RK, Tandon N, Kochupillai N. Prevalence and significance of low 25-hydroxyvitamin D concentrations in healthy subjects in Delhi. *The American journal of clinical nutrition*. 2000;72(2):472-5.
- Azizi F, RAIS ZF, Mir SGA. Vitamin D deficiency in a group of Tehran population. 2000.
- Sedrani SH. Low 25-hydroxyvitamin D and normal serum calcium concentrations in Saudi Arabia: Riyadh region. *Annals of nutrition and metabolism*. 1984;28(3):181-5.
- Alagöl F, Shihadeh Y, Boztepe H, Tanakol R, Yarman S, Azizlerli H, et al. Sunlight exposure and vitamin D deficiency in Turkish women. *Journal of endocrinological investigation*. 2000;23(3):173-7.
- Mansoor S, Habib A, Ghani F, Fatmi Z, Badruddin S, Mansoor S, et al. Prevalence and significance of vitamin D deficiency and insufficiency among apparently healthy adults. *Clinical biochemistry*. 2010;43(18):1431-5.
- Rab S, Baseer A. Occult osteomalacia amongst healthy and pregnant women in Pakistan. *The Lancet*. 1976;308(7997):1211-3.
- Zuberi LM, Habib A, Haque N, Jabbar A. Vitamin D deficiency in ambulatory patients. *Journal of the Pakistan Medical Association*. 2008;58(9):482.
- Atiq M, Suria A, Nizami S, Ahmed I. Maternal vitamin-D deficiency in Pakistan. *Acta obstetrica et gynecologica Scandinavica*. 1998;77(10):970-3.
- Mithal A, Wahl D, Bonjour J, Burckhardt P, Dawson-Hughes B, Eisman J, et al. IOF Committee of Scientific Advisors (CSA) Nutrition Working Group (2009) Global vitamin D status and determinants of hypovitaminosis D. *Osteoporosis Int*. 20:18071820.
- Nair R, Maseeh A. Vitamin D: The "sunshine" vitamin. *Journal of pharmacology & pharmacotherapeutics*. 2012;3(2):118.
- Riaz H, Finlayson A, Bashir S, Hussain S, Mahmood S, Malik F, et al. Prevalence of Vitamin D deficiency in Pakistan and implications for the future. *Expert review of clinical pharmacology*. 2016;9(2):329-38.
- Chaudhary B. Vitamin D deficiency in Rawalpindi-Islamabad region. *Journal of Rawalpindi Medical College*. 2017;21(2):169-72.
- Azizi S, Tariq TM. Vitamin D deficiency among Afghan adolescents in Kabul. *Journal of the College of Physicians and Surgeons Pakistan*. 2019;29(11):1072-7.
- Zargar A, Ahmad S, Masoodi S, Wani A, Bashir M, Laway B, et al. Vitamin D status in apparently healthy adults in Kashmir Valley of Indian subcontinent. *Postgraduate medical journal*. 2007;83(985):713-6.
- Rabbani A, Alavian S-M, Motlagh ME, Ashtiani MT, Ardalan G, Salavati A, et al. Vitamin D insufficiency among children and adolescents living in Tehran, Iran. *Journal of tropical pediatrics*. 2009;55(3):189-91.
- Tabrizi R, Moosazadeh M, Akbari M, Dabbaghmanesh MH, Mohamadkhani M, Asemi Z, et al. High prevalence of vitamin D deficiency among Iranian population: a systematic review and meta-analysis. *Iranian journal of medical sciences*. 2018;43(2):125.
- Atli T, Gullu S, Uysal A, Erdogan G. The prevalence of vitamin D deficiency and effects of ultraviolet light on vitamin D levels in elderly Turkish population. *Archives of gerontology and geriatrics*. 2005;40(1):53-60.
- Goswami R, Kochupillai N, Gupta N, Goswami D, Singh N, Dudha A. Presence of 25 (OH) D deficiency in a rural North Indian village despite abundant sunshine. *JAPI*. 2008;56:755-7.
- Mahmood K, Akhtar ST, Talib A, Haider I. Vitamin-D status in a population of healthy adults in Pakistan. *Methodology*. 2007.
- Al-Mutairi N, Issa B, Nair V. Photoprotection and vitamin D status: A study on awareness, knowledge and attitude towards sun protection in general population from Kuwait, and its relation with vitamin D levels. *Indian journal of dermatology, venereology and leprology*. 2012;78(3):342.
- Hagenau T, Vest R, Gissel T, Poulsen C, Erlandsen M, Mosekilde L, et al. Global vitamin D levels in relation to age, gender, skin pigmentation and latitude: an ecologic meta-regression analysis. *Osteoporosis international*. 2009;20(1):133-40.
- Sedrani SH, Elidrissi A, El Arabi KM. Sunlight and vitamin D status in normal Saudi subjects. *The American journal of clinical nutrition*. 1983;38(1):129-32.
- Looker AC, Pfeiffer CM, Lacher DA, Schleicher RL, Picciano MF, Yetley EA. Serum 25-hydroxyvitamin D status of the US population: 1988–1994 compared with 2000–2004. *The American journal of clinical nutrition*. 2008;88(6):1519-27.
- Looker AC, Dawson-Hughes B, Calvo M, Gunter E, Sahyoun N. Serum 25-hydroxyvitamin D status of adolescents and adults in two seasonal subpopulations from NHANES III. *Bone*. 2002;30(5):771-7.
- Shamim Khan A, Zubair M, Khan S, Sultan K, Khan MK. The role of vitamin D deficiency and supplementation in onset and progression of diabetic nephropathy: A systematic review. *Romanian Journal of Diabetes Nutrition and Metabolic Diseases*. 2022;29(1):135-54.
- Manicourt D-H, Devogelaer J-P. Urban tropospheric ozone increases the prevalence of vitamin D deficiency among Belgian postmenopausal women with outdoor activities during summer. *The Journal of Clinical Endocrinology & Metabolism*. 2008;93(10):3893-9.
- Hashemipour S, Larjani B, Adibi H, Javadi E, Sedaghat M, Pajouhi M, et al. Vitamin D deficiency and causative factors in the population of Tehran. *BMC Public health*. 2004;4(1):1-6.
- Gannagé-Yared MH, Chemali R, Yaacoub N, Halaby G. Hypovitaminosis D in a sunny country: relation to lifestyle and bone markers. *Journal of bone and mineral research*. 2000;15(9):1856-62.
- Moussavi M, Heidarpour R, Aminorroaya A, Pournaghshband Z, Amini M. Prevalence of vitamin D deficiency in Isfahani high school students in 2004. *Hormone Research in Paediatrics*. 2005;64(3):144-8.